

SH 365


.V7 M7

Copy 1



Class SH365

Book V7M7




DEPARTMENT OF COMMERCE AND LABOR
BUREAU OF FISHERIES
GEORGE M. BOWERS, Commissioner

CONDITION AND EXTENT OF THE
OYSTER BEDS OF JAMES
RIVER, VIRGINIA

Bureau of Fisheries Document No. 729



WASHINGTON
GOVERNMENT PRINTING OFFICE
1910



2

DEPARTMENT OF COMMERCE AND LABOR

BUREAU OF FISHERIES

GEORGE M. BOWERS, Commissioner

CONDITION AND EXTENT OF THE
OYSTER BEDS OF JAMES
RIVER, VIRGINIA

364
916

by H. F. Moore

Bureau of Fisheries Document No. 729



WASHINGTON
GOVERNMENT PRINTING OFFICE

1910

SH365
V9M7

JAN 22 1910
L. OF D.

31

03925

**CONDITION AND EXTENT OF THE OYSTER BEDS OF
JAMES RIVER, VIRGINIA**

BY H. F. MOORE

Assistant, U. S. Bureau of Fisheries

Bureau of Fisheries Document No. 729

P R E F A C E .

On February 3, 1909, the Bureau of Fisheries received from Hon. Claude A. Swanson, governor of Virginia, a communication inclosing the following resolution of the Commissioners of Fisheries of the State:


Resolved, That the governor be requested to enlist the services of the United States Bureau of Fisheries in determining and defining the fertile and the barren areas in James River, marking and platting same, provided it can be done without expenditure by the State.

At the urgent solicitation of Governor Swanson, and upon the conviction that the work would prove of value as a guide for contemplated legislation by the State in respect to the future administration of the public oyster grounds, the request for the survey was acceded to, the steamer *Fish Hawk* and civilian assistants were detailed for the work, and Dr. H. F. Moore, assistant in the Bureau of Fisheries, was directed to assume charge.

The erection of signals was begun early in July and completed by August 7. The actual examination of the oyster beds commenced on August 9 and extended, with only such interruptions as were due to the weather, to September 14, the survey thus covering the period just prior to the opening of the oyster season, when the beds were in their optimum condition. Under the terms of the resolution quoted above, the Bureau has not felt justified in offering advice as to the future treatment of the beds, and the following report is therefore confined to statements of fact and a short discussion of their several obvious avenues of application.

GEORGE M. BOWERS,
Commissioner.

UNITED STATES BUREAU OF FISHERIES,
Washington, D. C., December 1, 1909.



CONTENTS.

	Page.
Previous surveys.....	7
Methods of the present survey.....	9
Oyster rocks.....	13
Market oyster area—	
Hollands.....	14
Nansemond Ridge.....	15
Larkins.....	17
Drum Shoal.....	18
Newport News.....	19
Cruiser Shoal.....	20
Flat Rock and adjacent small beds.....	21
High Shoal.....	22
Trout Shoal.....	23
Dog Shoal.....	25
Fishing Point.....	26
Between Fishing Point and Ballards Marsh.....	27
Ballards Marsh.....	28
Creek Channel.....	29
Aaron Shoal.....	30
Browns Shoal.....	31
Gun.....	33
Kettle Hole.....	33
Thomas Point.....	35
Blunt Point.....	37
White Shoal.....	38
Seed oyster area—	
Jail Island.....	39
Wreck Shoal.....	41
Dry Shoals.....	42
Point of Shoals.....	43
Swash.....	45
Mulberry Swash.....	46
Marshy Island.....	47
Long Shoal.....	48
V Rock.....	50
Moores.....	51
Horsehead.....	51
Deepwater Shoals.....	53
Rock Wharf Shoals.....	54
Beds between Rock Wharf Shoals and Spindle Rock.....	54
Spindle.....	55
Days Point Shoal.....	56

	Page.
Public grounds.....	56
No. 2 Nansemond County and No. 6 Isle of Wight County.....	60
No. 1 Warwick County below Deep Creek.....	64
Minor public grounds.....	67
No. 1 Warwick County above Deep Creek.....	67
No. 1 Isle of Wight County.....	70
Summary.....	72
Market oyster area.....	72
Seed oyster area.....	77
Conclusion.....	80
Description of charts.....	83

CONDITION AND EXTENT OF THE OYSTER BEDS OF JAMES RIVER, VIRGINIA.

By H. F. MOORE,
Assistant, U. S. Bureau of Fisheries.

PREVIOUS SURVEYS.

Prior to the investigations made by the Bureau of Fisheries in July, August, and September, 1909, two surveys of the James River oyster beds had been made, neither of which professed to delineate the rocks accurately or to furnish detailed information concerning their productiveness and condition. The first of these surveys was a reconnoissance made in 1878 by Lieut. (then Master) Francis Winslow, U. S. Navy, in command of the Coast and Geodetic Survey schooner *Palinurus*. The second was the survey of the public grounds by Mr. J. B. Baylor, assistant, Coast and Geodetic Survey, under the authority of the State, in 1892 and preceding years.

As Winslow himself states, his "examination of these beds was a very hurried one, and the delineation must be regarded as merely approximate, being the result of a hasty reconnoissance." The chart published with the report delineates merely the general outlines of the oyster-bearing areas, without attempting to show the smaller individual rocks or the density of growth, and the text is of very general character. Comparing the chart with the results of the recent survey, however, it is evident that Lieutenant Winslow's brief investigation must have shown with considerable accuracy the general distribution of oysters in the James and Nansemond rivers at that time. The differences between the general results of the two surveys are such as could be readily produced by the lapse of time and the vicissitudes through which the beds have passed under the operation of natural and human agencies. Some areas have become depleted through the intensive fishing they have sustained, or from the effects of freshets and other physical factors, while on the other hand some appear to have had their boundaries extended or have become merged with adjacent beds through the operations of the tongers.

The Baylor survey was of an entirely different character from that conducted by Winslow. It was in no sense an examination of the oyster beds themselves, but primarily and avowedly a delimitation of boundaries which included the recognized or reputed oyster-bearing bottom, as pointed out by local commissioners or representatives of each oyster-producing county. It is the writer's understanding that the county commissioners were, under the state law ordering the survey, the final arbiters with whom rested the decision whether or not a given area should be included within the boundaries of the public grounds. So far as can be learned no examinations whatever were made on the beds, the commissioners using their judgment and local knowledge in selecting the corners and the engineers with their theodolites cutting in the points indicated from stations on the shore.

Whether or not beds were omitted from the confines of the public grounds so located can not now be satisfactorily determined, owing to the development of the planting industry, outside of the Baylor lines, on all or most of the available bottom. It is evident, however, that in the region under discussion no very extensive rocks were disregarded, and a comparison of the results of the recent survey with that of 1892 shows that the Baylor lines, considered as a broad scheme of delimitation, conform closely with the general distribution of the rocks. At several places, notably on Gun and Kettle Hole rocks, parts of the natural beds undoubtedly fell outside of the lines, but the writer hazards the suggestion that this may not have been through inadvertence but because those parts of the rocks may have been already taken up as private holdings.

It has been claimed, and Mr. Baylor himself has so stated in official communications to the State, that a very considerable area of barren bottom, amounting to many thousand acres, was included within the public grounds. That this should be so, under the system adopted by the local commissioners and under the desire to assure the inclusion of all naturally productive bottom, was inevitable. Moreover, the boundaries of the beds are irregular curves, while the including surveyed boundaries must be straight lines, for purposes of administration and policing as long and unbroken as possible. To have excluded the greater part of this barren bottom would have necessitated a careful location of the natural rocks and the breaking up of the public grounds into a considerable number of small or moderate areas instead of segregating them into a few large ones.

To what extent the claim that great areas of barren bottom are included in the public grounds is justified will appear from the accompanying charts and in the following descriptions and discussions.

METHODS OF THE PRESENT SURVEY.

To furnish authoritative and definite information as to the actual extent and condition of the natural rocks and the character of the bottoms embraced within the boundaries of the public beds, it was necessary to depart widely from the methods of the previous surveys.

It was decided to confine the investigation wholly to the public beds, passing their boundaries only far enough to give assurance that the entire area had been covered. Nothing was to be gained by an examination of the excluded areas, as it is now almost impossible to determine whether natural rocks were omitted from the grounds laid out in 1892, and it is too late to correct such omissions if they could be determined. For legal purposes, all that is not avowedly public ground is barren bottom, and if held under leasehold from the State can not be alienated from the possession of the lessees as long as the law has been complied with.

The methods followed have been essentially those pursued in former surveys conducted by the Bureau of Fisheries, with the changes and improvements dictated by recent experience and the local conditions.

The Coast and Geodetic Survey furnished projections on which were platted the triangulation points used in former surveys by that bureau. Several of these points, including the light-houses, were "recovered," and from them the signals, usually tripods, erected where necessary, were cut in and platted by means of the sextant and 3-arm protractor. This method, while lacking the great precision attained by means of the best theodolites and the nice computations employed by the Coast Survey in its work, insures an accuracy more than sufficient for the purposes of an oyster survey.

The oyster beds were discovered by soundings with a lead line, but principally by means of a length of chain dragged over the bottom at the end of a copper wire running from the sounding boat. The wire was wound on a reel and its unwound length was adjusted to the depth of water and the speed of the launch, so that the chain was always on the bottom. Whenever the chain touched a shell or an oyster the shock or vibration was transmitted up the wire to the hand of a man whose sole duty it was to give heed to such signals and report them to the recorder.

The launches from which the soundings were made were run at a speed of between 3 and 4 miles per hour, usually on ranges ashore to insure the rectitude of the lines. At intervals of three minutes—in some cases two minutes—the position of the boat was determined by two simultaneous sextant observations of the angles between a set of three signals, the middle one of which was common to the two angles, the position being immediately platted on the boat sheet. At regular intervals of twenty seconds, as measured by

a clock under the observation of the recorder, the leadsman made a sounding and reported to the recorder the depth of water and the character of the bottom, immediately after which the man at the wire reported the character of the chain indications since the last sounding—that is, whether they showed barren bottom or dense, scattering, or very scattering growths of oysters.

With the boat running at 3 miles per hour the soundings were between 80 and 90 feet apart and, as the speed of the boat was uniform, the location of each was determinable within a yard or two by dividing the platted distance between the positions determined by the sextant by the number of soundings. The chain, of course, gave a continuous indication of the character of the bottom, but the record was made at the regular twenty-second intervals observed in sounding.

The chain, while indicating the absence or the relative abundance of objects on the bottom, gives no information as to whether they are shells or oysters, nor, if the latter, their size and condition. To obtain this data it was necessary to supplement the observations already described by others more definite in respect to the desired particulars. Whenever in the opinion of the officer in charge of the sounding boat such information was required, a numbered buoy was dropped, the time and number being entered in the sounding book. Another launch, following the sounding boat, anchored alongside the buoy, and a quantity of the oysters and shells were tonged up, separated by sizes, and counted.

In former surveys made by the writer, in order to arrive at an estimate of the density of the oyster growth a definite area, usually 5 yards, was staked off by means of steel-shod pikes and everything was removed from the bottom and counted. This method is accurate, but slow and difficult in deep water, and, as it was desirable to make a large number of observations, the system developed in the Maryland survey was adopted. This consists essentially in making a known number of "grabs" with the oyster tongs, exercising care to clean the bottom of oysters as thoroughly as possible at each grab. In a given depth of water and using the same boat and tongs an oysterman will cover practically the same area of the bottom at each grab, but, other factors remaining the same, the area of the grab will decrease with an increase in the depth.

Careful measurements were made and tabulated showing the area per grab covered by the tonger employed on the work at each foot of depth of water and for each pair of tongs and boat used. With this data, and knowing the number of "grabs," the number of oysters of each size per square yard of bottom was readily obtainable by simple calculation. The following example will illustrate the data obtained and the form of the record:

DEPARTMENT OF COMMERCE AND LABOR.

BUREAU OF FISHERIES.

*Field record of examinations of oyster beds.*General locality: *James River.*Local name of oyster ground: *Between Rock wharf and Spindle rock.*Date: *August 28, 1909.*Time: *9.25 a. m.*Angle: *F. 140.*Buoy No.: *23.*Depth: *7 feet.*Bottom: *Hard.*Condition of water: *Medium clear.*

Density: Temperature:

Current: Stage of tide:

Tongman: *Lawrence, in flatboat.*No. grabs made: *8.*Tongs: *14 feet.*Total area covered: *3½ square yards*No. oysters taken: { -1 in.: *27.* 1 in.—X in.: *69.*X in.—4 in.: *10.* 4 in.: *3.*Quantity shells: *½ bushel.*Result: { Spat per square yard: *7.7.*
Culls per square yard: *19.7.*
Counts per square yard: *3.7.*

X in. = cull limit prescribed by law.

This furnishes an exact statement of the condition of the bed at a spot which can be platted on the chart with error in position of not more than a few yards. From the data obtained a close estimate may be formed of the bushels of oysters and shells per acre in the vicinity of the examination and, by multiplying the observations, for the bed as a whole. In the course of the survey 590 observations were made at various places, principally on the natural rocks, but some on the barren bottoms also.

In former surveys by the Bureau the relative density of the oyster growth has been considered solely from the standpoint of the total quantity of oysters. That method is satisfactory where the depth is fairly uniform throughout the region examined, but was not considered accurate enough for the purposes of the present report.

With a given quantity of oysters per square yard or acre, a bed lying in shoal water is more valuable commercially than one in deep water, owing to the fact that the labor of the tonger is more efficient in the former. As has been pointed out, the area covered by a "grab" decreases with an increase in depth, and, moreover, the deeper the water the greater is the labor involved in making the "grab" and the smaller is the number of grabs which can be made per hour or per day. With 14-foot tongs used from a canoe, such as is employed

on the James River, an oysterman can cover twice as much bottom per grab in 4 feet as he can in 8 feet, and about two and one-half times as much as he can using 20-foot tongs in 16 feet. Using the tongs stated, the average tonger observed in Maryland, and the data will hold in Virginia, will make about 2.7 grabs per minute in 4 feet, 2.6 in 8 feet, and 1.8 in 16 feet of water. In other words, if he can cover 1 square yard of the bottom in a given time in 16 feet, he can cover 1.7 yards in 8 feet, and 3.3 yards in 4 feet of water.

It is obvious that if a tonger in a given time is to obtain the same quantity of oysters in each of these depths, the oyster growth must lie on the bottom with a density inversely to the areas stated above. The value of a bed, the price per bushel of the oysters being the same, depends on the quantity which a man can take in a given time, and it therefore happens that a bed in deep water may be valueless commercially, while another rock, with the same density of growth but covered by shoaler water, may be tonged with profit.

Based on these principles, and taking into consideration the number of oysters per bushel on the different beds as determined by actual counts, tables were prepared showing the number of oysters per square yard for each foot of depth necessary to yield to the tonger 1 bushel of oysters per day of tonging. From these data the beds were divided into areas, according to the number of bushels of oysters which they were capable of yielding per day to the tonger, based on nine hours of actual tonging and disregarding the time occupied in culling. The bottom was divided into 5 categories: Barren, on which there were neither shells nor oysters; depleted, on which the tonger could take less than 3 bushels of market oysters or 4 bushels of seed, according to location; very scattering growth, on which between 3 and 5 bushels of oysters or 4 and 8 bushels of seed could be taken; scattering growth, on which the limits were 5 and 8 bushels of market oysters or 8 and 12 bushels of seed; and areas of dense growth, on which upward of 8 bushels of market oysters or 12 bushels of seed could be taken per day.

During the survey 10,440 soundings were taken, and the position of the boat was instrumentally determined at 1,369 places. The chain was dragged for 226 miles, giving continuous indications of the character of the bottom, which were plotted on the chart at 10,440 places. The density of oyster growth was determined by the 590 biological observations already referred to, and the extent and boundaries of the areas as charted were fixed by a combination of these observations and the 10,440 records of the continuous chain readings. During the work the writer was in charge of the sounding boat and in constant touch with all operations. The biological observations were all under the immediate charge of Mr. T. E. B. Pope, whose

experience in such work is such as insured their accuracy both as to the area covered by the tonger and the quantity of oysters taken. The oysters brought in by the biological party were all examined by the author, who has also personally made all of the many calculations required and directly supervised the laying off of the areas on the charts. The basis for the determination of the character of the beds was decided on in advance, but the work of the survey was so planned that it was impossible for any member of the party to form an opinion as to the conditions found until after the field work was completed, and any involuntary prejudice was thus eliminated as far as possible. The author himself could form but a vague idea of the general results until the charts were completed and the report almost written.

In the following pages the subject is gradually developed from a detailed description of the several parts of the individual natural rocks to a broad consideration of the market oyster and seed areas as a whole, and in every case there is given the principal data on which the several statements are based.

OYSTER ROCKS.

The term "oyster rock," as used in Virginia and employed in this report, is synonymous with natural oyster bed and is to be distinguished from the term "public ground," which is used to designate the areas legally embraced within surveyed lines and set apart for the use of the public. The public grounds were intended to embrace all of the oyster rocks, and usually each includes a number of the latter within its confines.

An oyster rock is usually a more or less definite area of bottom, limited by the extent of actual oyster growth. Originally, the boundaries were rather definitely marked and the rocks were separated from one another by barren areas, but the operations of oystering have in many cases strewn oysters and shells over the surrounding bottom, so that in cases the original limits have become obscured and adjacent rocks merged.

On the accompanying chart much of the bottom indicated as depleted really represents the areas which have been thus covered with scattered oysters and shells, and the term employed indicates that oysters and shells are very scarce rather than that they have been removed, though the latter is the fact in many cases. The so-called "depleted" areas are those on which oysters grow in quantities much below those which would make it commercially profitable to tong for them.

The boundaries of the rocks, as shown by the red inclosing lines on the charts and as considered in the text, were defined by the

results of the chain indications, the methods of obtaining which have been before explained. All areas in which shells or oysters were encountered are regarded as rocks, but their character, so far as productiveness is concerned, was determined by tonging and counting the yield in the manner heretofore described. The depleted areas, except where it is shown that they contain a reasonably heavy growth of young, may be regarded as worthless from the viewpoint of the tonger; the areas of very scattering growth are of doubtful value except where a heavy growth of young oysters indicates potential improvement, while the areas of scattering and dense growth can be regarded as really productive natural rocks. The barren bottom, which is shown on the chart as an unshaded area outside of the red lines, is that on which neither oysters nor shells were found. A few small unshaded areas inclosed by red lines indicate beds the exact nature of which was not accurately determined.

An attempt is made in this report to designate the rocks by the names employed by the oystermen, so far as these could be ascertained. In several cases, as for instance "Fishing Point Rocks" and "Marshy Island Rock," names have been coined to serve the purposes of reference and designation. The exact extent of Point of Shoals Rock was not definitely ascertained, and as shown on the chart it may not accord with the usage of the oystermen. There was also some doubt about the location of Kettle Hole and Thomas Point Rocks, but, as the names used in the text are clearly shown on the charts, there can be no confusion in the references. In a number of cases where the several beds were more or less continuous with one another arbitrary boundaries have been adopted, but, as these usually pass through depleted areas and as in a later discussion the rocks are considered as a whole in their grouping in the public beds, the necessity for this treatment causes no loss in the final accuracy or exactitude.

In the following pages the rocks are considered in detail.

MARKET OYSTER AREA.

HOLLANDS ROCK.

This was intended to be included by the Baylor survey in Public Ground No. 3, Nansemond County, though it is stated that a mistake was made by which it was omitted. The area, 22 acres, which is described under this name included the only bottom within the public ground which gave any indication of being an oyster bed, though the examination showed it to be depleted. It is completely surrounded by planted beds.

The results of the examination were as follows:

DETAILS OF EXAMINATION OF HOLLANDS ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity of oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
4 30	Aug. 9, 1909	Feet. 11.0	Depleted.....	0	0.4	1.8	<i>Bush.</i> 3	<i>Bush.</i> 29	<i>Bush.</i> 32
	Aug. 11, 1909	9.5do.....	0	0	0	0	0	0

NANSEMOND RIDGE ROCK.

This is the principal and only productive bed in Nansemond River. It lies mainly on and about a shoal extending through the middle of the river from Cedar Point almost to the middle of James River, opposite Newport News. At its northern end it is connected, by an unproductive, practically barren area, with three smaller rocks hereafter described. Its area, density of oyster growth, and contents are as follows:

OYSTER GROWTH ON NANSEMOND RIDGE ROCK.

Character of growth of market oysters.	Area.	Oysters per acre.		Estimated content of market oysters.
		Seed.	Market.	
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	85	148	93	7,905
Scattering.....	446	133	60	26,760
Very scattering.....	294	34	37	10,878
Depleted.....	782	35	11	8,602
Total.....	1,607			54,145

The market oysters on this bed are large, averaging at the time of the survey a few over 300 per bushel. They are said to attain a good condition, particularly late in the season, and are used mainly by shucking houses. The small oysters ran about 750 per bushel.

The broadest, largest, and most productive part of the bed stretches northward from opposite Pig Point on the west side of the channel. It is estimated that this portion has an area of about 1,156 acres, of which 69 acres bear a dense growth of market oysters, 386 acres a scattering growth, 201 acres a very scattering growth, and 500 acres are depleted. The latter does not include the barren bottom embraced between the edges of the bed and the lines of the Baylor survey. It is further estimated that on the dense bottom a man tonging exclusively could take in a day about 10 bushels of market oysters, on the scattering area about 6 bushels, on the very scattering part about 3½ bushels, while on the depleted area he could not take an average of over 1 bushel. These estimates are for the beginning of the season,

and any considerable tonging of the beds would soon materially reduce the average catch per day.

In the dense and scattering parts of this portion of the bed, especially near the crest of the ridge, there is a growth of small oysters so dense that an average of upward of 12 bushels could be tonged per day, and these areas can undoubtedly be regarded as both presently and prospectively productive. There is also a dense growth of young oysters on the inner parts of the depleted area opposite Nansemond River Light. On the areas of very scattering growth the small oysters are in even smaller quantity than the market oysters, but in places there are clean shells in sufficient quantity to indicate that under proper conditions a good set might occur and the bottom become fairly productive.

Above a line drawn between Pig Point and Barrel Point the bed may be divided into two parts, one a tail-like continuation of the main bed running along the eastern edge of the channel and the other a detached portion lying on a shoal west of the channel, north of Lar-kins Rock. The former has 126 acres of depleted bottom and two small patches, one of scattering growth covering about 22 acres and the other of about 41 acres on which the oysters are very scattering. The detached area covers about 260 acres, of which 15 are dense, 39 scattering, 51 very scattering, and 155 depleted. On the areas of dense and scattering growths of market oysters there is a heavy growth of culls, but the scattering and depleted areas are generally impoverished of young.

On the two areas just described as lying above a line between Pig and Barrel points it is estimated that there are 15 acres of dense growth on which a man could tong an average of about 8 barrels of market oysters per day, 61 acres on which he could average about 5 bushels, 92 acres of very scattering growth where he could take about 4 bushels per day, and 281 acres of depleted bottom which will not yield 1 bushel per day. On the depleted area there are few young oysters and practically no shells. The barren bottom lying within this part of the Baylor survey, on which oysters do not now grow and apparently never have grown in marketable quantities, nearly equals all of the foregoing combined, covering about 430 acres. The barren and depleted bottom together aggregate about 711 acres, while all of the bottom which is capable of yielding even as little as 3 bushels per day, exclusive of the time consumed in culling, covers about 168 acres. In other words, at least 80 per cent of the area is at present commercially worthless.

The observations, in addition to the sounding and chain investigations, on which the foregoing is based, are as follows:

DETAILS OF EXAMINATION OF NANSEMOND RIDGE ROCKS.

Station number.	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
		<i>Feet.</i>					<i>Bush.</i>	<i>Bush.</i>	<i>Bush.</i>
1	Aug. 9, 1909	13.0	Dense.....	6.0	5.1	8.2	72	132	204
549	Sept. 13, 1909	6.5	do.....	4.7	27.7	4.7	209	76	285
582	Sept. 14, 1909	6.5	do.....	3.9	21.2	4.4	163	71	234
15	Aug. 10, 1909	6.0	Scattering.....	1.7	13.4	2.7	98	43	141
18	do.....	11.5	do.....	.2	.0	4.0	1	64	65
33	Aug. 12, 1909	10.5	do.....	.6	12.4	4.6	85	74	159
34	do.....	7.5	do.....	4.8	13.3	3.6	111	58	169
42	do.....	10.5	do.....	.2	.6	4.5	5	72	77
550	Sept. 13, 1909	6.5	do.....	2.3	11.6	2.3	90	37	127
576	Sept. 14, 1909	7.5	do.....	1.8	9.7	3.1	75	50	125
577	do.....	8.0	do.....	45.0	26.6	3.9	466	63	529
578	do.....	11.0	do.....	4.0	16.3	4.0	132	64	196
579	do.....	6.0	do.....	5.3	27.3	3.5	212	56	268
580	do.....	7.5	do.....	5.3	25.6	4.2	201	68	269
581	do.....	8.5	do.....	3.4	15.7	4.9	124	79	203
7	Aug. 9, 1909	10.0	Very scattering.....	2.9	9.3	1.8	79	29	108
21	Aug. 10, 1909	9.0	do.....	.4	2.1	2.7	16	43	59
26	Aug. 11, 1909	8.5	do.....	1.6	10.9	2.8	81	45	126
35	Aug. 12, 1909	9.5	do.....	.9	3.8	2.2	31	35	66
39	do.....	10.0	do.....	.2	.4	2.3	4	37	41
43	do.....	10.5	do.....	.8	2.4	3.1	21	50	71
542	Sept. 13, 1909	6.0	do.....	.0	.8	1.4	5	23	25
2	Aug. 9, 1909	12.0	Depleted.....	.0	.0	.9	0	15	15
5	do.....	9.0	do.....	3.5	20.0	.9	153	15	168
8	do.....	8.0	do.....	8.4	23.9	.6	210	10	220
11	Aug. 10, 1909	8.5	do.....	1.8	13.3	.5	98	8	106
12	do.....	7.0	do.....	2.1	4.5	.9	43	14	57
13	do.....	7.5	do.....	.0	.0	.3	0	5	5
16	do.....	12.5	do.....	.0	.0	.4	0	6	6
25	Aug. 11, 1909	10.0	do.....	.0	.0	.0	0	0	0
36	Aug. 12, 1909	10.5	do.....	.5	1.8	1.3	8	21	29
38	do.....	9.0	do.....	.7	.9	1.3	10	21	31
40	do.....	10.0	do.....	.4	.7	1.2	7	19	26
41	do.....	10.0	do.....	.7	3.5	1.9	27	31	58
547	Sept. 13, 1909	7.0	do.....	.2	.7	1.1	6	18	24
548	do.....	6.5	do.....	.0	.0	.2	0	3	3
551	do.....	7.5	do.....	1.3	2.9	.5	27	8	35
574	Sept. 14, 1909	9.5	do.....	.0	.0	.0	0	0	0
575	do.....	9.0	do.....	.0	.0	.0	0	0	0
583	do.....	7.0	do.....	2.6	10.0	1.1	82	18	100
584	do.....	10.5	do.....	.0	.0	.0	0	0	0
587	do.....	11.0	do.....	.0	.0	.0	0	0	0

LARKINS ROCK.

This is a small bed in Nansemond River at the extreme southwest corner of Public Ground No. 2. As developed by the survey it has an area of about 39 acres and a depth varying from 4½ to 8 feet at mean-low water. It is stated that the product of this bed has been in demand by shucking houses, the size and quality being generally good and the condition fat, especially early in spring. The market oysters found by the survey averaged between 300 and 350 to the bushel and the small oysters about 750 per bushel.

The bed at present bears market oysters at the average density of about 5 bushels per acre, though in spots the production is as high as 18 bushels. The young growth has an average density of about 10 bushels and a maximum of 31 bushels per acre.

At the present time this bed must be regarded as depleted, as at none of the spots examined could a man tong more than 2 bushels of oysters per day, and the average yield, taking the bed as a whole,

would be hardly more than one-half bushel per day. The young growth is sparse and the shells few. The bed bears the aspect of having been carried off bodily for planting purposes, a depredation to which its location makes it susceptible. The results of detailed examinations are as follows:

DETAILS OF EXAMINATION OF LARKINS ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
		<i>Fect.</i>					<i>Bush.</i>	<i>Bush.</i>	<i>Bush.</i>
20	Aug. 10, 1909	8.5	Depleted.....	0.0	0.0	0.0	0	0	0
27	Aug. 11, 1909	7.5	do.....	0	0	0	0	0	0
28	do.....	6.5	do.....	0	3.5	.1	23	2	25
544	Sept. 13, 1909	6.5	do.....	0	0	0	0	0	0
545	do.....	7.0	do.....	.2	2.7	1.1	31	18	49
546	do.....	6.5	do.....	.0	.6	.6	4	10	14

DRUM SHOAL ROCK.

This is a small bed located at the northwest corner of Public Ground No. 2 in Nansemond County. Its area, density of oyster growth, and estimated contents are as follows:

OYSTER GROWTH ON DRUM SHOAL ROCK.

Character of growth of market oysters.	Area.	Oysters per acre.		Estimated content of market oysters.
		Seed.	Market.	
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Scattering.....	19	92	50	950
Very scattering.....	14	61	39	546
Depleted.....	95	62	29	2,755
Total.....	128			4,251

This bed was doubtless originally restricted to the area of the shoal which is now covered by the scattering and very scattering growth, but oysters and shells have become scattered over the surrounding bottom and it is now connected, by means of a depleted area, with Nansemond Ridge Rock on the south and Newport News Rock on the west.

The scattering growth lies in a depth of from 6 to 8 feet at mean low water and the market oysters grow in such quantity that a tonger of average ability can take about 5 bushels per day. The very scattering growth is at the eastern side of the shoal and has oysters in sufficient numbers to yield about $3\frac{1}{2}$ bushels per day. The density of growth shown for the depleted area in the table produced above is in excess of the actual conditions, as the examina-

tions on which it was based were taken in close proximity to the edge of the shoal, while the more distant bottom is more denuded. The growth of young on the productive part of the rock is fair, being sufficient to yield about 9 bushels per day on the scattered area and about 6 bushels on the very scattered area and about the edges of the shoal.

Following is the record of observations on this bed:

DETAILS OF EXAMINATION OF DRUM SHOAL ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
585	Sept. 14, 1909	8.0	Scattering.....	2.6	11.6	3.1	<i>Bush.</i> 92	<i>Bush.</i> 50	<i>Bush.</i> 142
586	do.....	9.0	Very scattering....	1.5	7.9	2.4	61	39	100
44	Aug. 12, 1909	10.5	Depleted.....	.7	6.1	1.9	44	31	75
45	do.....	8.5	do.....	.9	12.0	1.7	84	27	111

NEWPORT NEWS ROCK.

This lies in the overlapping portions of Public Grounds No. 2, Nansemond County, and No. 6, Isle of Wight County, north of Nansemond Ridge, and between Drum Shoal on the east and Cruiser Rock on the west. Its estimated area, density of growth, and contents are as follows:

OYSTER GROWTH ON NEWPORT NEWS ROCK.

Character of growth of market oysters.	Area.	Oysters per acre.		Estimated content of market oysters.
		Seed.	Market.	
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	4	93	108	432
Scattering.....	27	75	63	1,701
Very scattering.....	12	83	35	420
Depleted.....	129	34	27	3,483
Total.....	172			6,036

The rock in reality consists of several shoal spots separated by areas of depleted bottom in deeper water. The dense area is a small spot lying by itself in about 8 feet of water at low tide, and it bears market oysters in sufficient quantity to yield to the tonger about 9 bushels per day, and the young growth is in nearly the same quantity. The scattered and very scattered growth lies on Cruiser Shoal proper, the former being sufficiently productive to yield about 6 bushels and the latter about 3 bushels per day. On both of these areas there is a fair growth of young, sufficient to yield about 6 bushels per day.

At the eastern edge of the rock, as defined on the chart, there is a dense growth, not shown, which lies just outside of the Baylor line, and running south from this is a growth of young oysters on the so-called depleted bottom sufficient to yield about 4 bushels per day. The depleted bottom on the whole will yield about 2 bushels of market oysters per day and about the same quantity of young.

The following are the results of examinations on this rock:

DETAILS OF EXAMINATION OF NEWPORT NEWS ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
		<i>Feet.</i>					<i>Bush.</i>	<i>Bush.</i>	<i>Bush.</i>
52	Aug. 13, 1909	12.0	Dense.....	0.7	13.6	6.7	93	108	201
53do.....	9.0	Scattering.....	1.1	9.8	4.7	71	76	147
55do.....	11.0do.....	1.8	10.2	3.1	78	50	128
56do.....	10.0	Very scattering.....	1.1	11.6	2.2	83	35	118
23	Aug. 11, 1909	9.5	Depleted.....	3.1	5.1	1.6	53	26	79
46	Aug. 12, 1909	9.5do.....	.7	4.9	1.8	36	30	66
54	Aug. 13, 1909	10.0do.....	1.1	1.1	1.6	14	26	40

CRUISER SHOAL ROCK.

This rock lies on and about the shoal that gives it its name, mainly in Public Ground No. 6, Warwick County, but partly in the area common to that ground and No. 2, Nansemond County. Its area and condition are shown in the following table:

OYSTER GROWTH ON CRUISER SHOAL ROCK.

Character of growth of market oysters.	Area.	Oysters per acre.		Estimated content of market oysters.
		Seed.	Market.	
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	27	140	156	4,212
Scattering.....	19	47	53	1,007
Very scattering.....	26	51	28	728
Depleted.....	32	66	9	288
Total.....	104			6,235

The dense and scattered areas follow the line of a very shallow ridge which forms the backbone of the shoal, the former being capable of yielding from 10 to 20 bushels of oysters per day and the latter about 5. The area of very scattering growth lies on each side of the more prolific areas and is capable of yielding about 3 bushels of market oysters per day's tonging. The depleted area will yield an average of not over 1 bushel of market oysters per day, and the parts more distant from the ridge are practically bare. Close to the ridge, even on some of the bottom depleted of market oysters, the

growth of young is good enough to yield an average of about 15 bushels per day to the tonger; but on the very scattering and depleted areas further removed from the ridge the young growth is sparse.

The following table shows the results of the examinations made on this bed:

DETAILS OF EXAMINATION OF CRUISER SHOAL ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
58	Aug. 13, 1909	<i>Feet.</i> 10.5	Dense.....	4.0	21.0	7.3	<i>Bush.</i> 162	<i>Bush.</i> 117	<i>Bush.</i> 279
589	Sept. 14, 1909	8.5	do.....	1.5	17.0	12.1	120	195	315
51	Aug. 12, 1909	9.0	Scattering.....	.3	7.0	3.3	47	53	100
50	do.....	11.0	Very scattering.....	.2	.4	2.0	4	32	36
57	Aug. 13, 1909	7.5	do.....	1.8	13.4	1.6	99	25	124
47	Aug. 12, 1909	9.0	Depleted.....	.5	4.5	.7	32	11	43
59	Aug. 13, 1909	13.0	do.....	.0	.0	.0	0	0	0
590	Sept. 14, 1909	6.0	do.....	5.3	20.2	.9	166	15	181

FLAT ROCK AND ADJACENT SMALL BEDS.

Flat Rock is a small bed bearing a dense growth of market oysters lying in the southeast corner of Public Ground No. 6, Warwick County. The examination of this rock was not satisfactory, as owing to an error in platting in the field certain positions supposed to be on the bed proved to be on adjacent planted beds. The single examination, in connection with traversing lines of chain readings, indicates a growth over the entire area which will yield to the tonger an average of about 9 bushels of market oysters per day. There were practically no small oysters or shells, and there was some reason to believe that the place had been planted, though it was fully 200 yards inside of the Baylor lines.

North of Flat Rock is a small depleted area, covering about 7 acres, on which there are about 26 bushels of market oysters and 16 bushels of young oysters per acre, and on which it is computed that a tonger could take not over 2 bushels of oysters per day.

West of Cruiser Rock is another unnamed bed of very scattering oysters. Its area is about 5 acres, with an average of 42 bushels of market oysters and 55 bushels of seed oysters per acre, and it is estimated that a tonger could take about 3½ bushels of oysters per day.

Northwest of Flat Rock, at intervals of about 400 yards, are two small beds where the water does not shoal, on which no determinations were made except with the chain. The indications are of very scattering growths. The areas are about 5 and 2 acres, respectively. The data relating to the several beds examined are as follows:

DETAILS OF EXAMINATION OF FLAT ROCK AND SMALL BEDS BETWEEN NANSEMOND
RIDGE AND FISHING POINT.

Station number.	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
		<i>Feet.</i>					<i>Bush.</i>	<i>Bush.</i>	<i>Bush.</i>
31	Aug. 11, 1909	10.5	Dense.....	0.2	0.2	6.7	3	108	111
48	Aug. 12, 1909	10.0	Very scattering.....	1.8	6.7	2.6	55	42	97
32	Aug. 11, 1909	10.5	Depleted.....	.0	2.4	1.6	16	26	42

HIGH SHOAL ROCK.

High Shoal Rock is conspicuous from its position, near the middle of James River, surrounding a shoal of sand and broken shells bare at practically all times. The highest part of the shoal is near the channel, from which it extends shoreward toward Fishing Point. The bed, including the depleted parts, is quadrangular in shape, with its more productive areas extending at right angles to the shores along its major diameter. Its extent and density of growth are as follows:

OYSTER GROWTH ON HIGH SHOAL ROCK.

Character of growth of market oysters.	Area.	Oysters per acre.		Estimated content of market oysters.
		Seed.	Market.	
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	24	134	90	2,160
Scattering.....	13	127	48	624
Very scattering.....	24	58	25	600
Depleted.....	95	8	8½	807
Total.....	156			4,191

The dense area extends along practically the entire length of the shoal, as a narrow strip on both sides but especially to the eastward of the highest ridge. The market oysters are somewhat smaller than those in water a little deeper, but on the bed as a whole they were found to average about 400 to the bushel. It is estimated that on this area a tonger could take about 10 bushels of oysters per day.

The area of scattering growth forms a strip on the southern side of the bed along the edge of a deep swash channel which separates it from one of the neighboring Fishing Point Rocks. Market oysters are produced in sufficient abundance to yield the tonger about 6½ bushels per day. The very scattering area lies to the eastward of the dense area and north of the scattering, and produces oysters sufficient to furnish the tonger about 3 to 3½ bushels per day.

The depleted area, which constitutes the major portion of the bed as delineated on the chart, is principally on the western or upstream side, though a strip averaging about 100 yards in width extends around the outer end of the shoal and along its entire eastern side. This area will nowhere yield to the tonger more than about 2½ bushels of oysters per day, and the average yield of all places examined would not be over 1 bushel.

The growth of young oysters on the areas charted as dense and scattering and on the very scattering part closer to the ridge is prolific enough to yield a tonger from 8 to 25 bushels per day, the average of all places examined being about 17 bushels. Clean shells were abundant on the three productive areas and in the depleted area close to the ridge.

The data on which the foregoing statements are based is as follows:

DETAILS OF EXAMINATION OF HIGH SHOAL ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
		<i>Fect.</i>					<i>Bush.</i>	<i>Bush.</i>	<i>Bush.</i>
68	Aug. 13, 1909	7.0	Dense	3.6	10.9	7.8	94	94	188
70	do	7.5	do	10.0	17.0	10.3	175	125	300
473	Sept. 7, 1909	5.5	do	3.2	12.4	2.6	101	32	132
474	do	6.0	do	3.2	22.5	8.4	167	102	269
67	Aug. 13, 1909	6.0	Scattering	3.4	26.4	4.1	194	50	244
478	Sept. 8, 1909	6.0	do	2.9	6.5	3.9	61	47	108
476	do	8.0	Very scattering	0	3	2.9	2	35	37
477	do	3.5	do	7.6	10.1	1.3	115	16	131
63	Aug. 13, 1909	14.5	Depleted	0	0	0	0	0	0
64	do	12.5	do	0	2	1.8	1	22	23
69	do	9.0	do	0	0	0	0	0	0
71	do	7.5	do	4	2.2	1.8	17	22	39
470	Sept. 7, 1909	4.0	do	2.4	1.8	2	27	2	29
471	do	6.0	do	1.1	1.4	1.7	16	20	36
472	do	6.5	do	0	1.0	1.7	6	20	26
479	Sept. 8, 1909	7.5	do	0	2	0	1	0	1
480	do	9.5	do	0	1.3	0	8	0	8
481	do	7.0	do	0	0	0	0	0	0

TROUT SHOAL ROCK.

This bed occupies the southeastern part of Naseway Shoal. In its depleted area it is continuous with Dog Shoal Rock, which occupies the upper part of the same shoal, but is separated from the adjacent Fishing Point and High Shoal Rocks by swash channels in which there is an abrupt deepening of the water. The depth ranges from low-water mark to 10 or 11 feet, the greater depths being found in a pocket of deep water which extends far into Naseway shoal from the westward.

The extent and productiveness of the bed are shown in the following table:

OYSTER GROWTH ON TROUT SHOAL ROCK.

Character of growth of market oysters.	Area.	Oysters per acre.		Estimated content of market oysters.
		Seed.	Market.	
	Acres.	Bushels.	Bushels.	Bushels.
Scattering.....	25	165	44	1,100
Very scattering.....	14	118	30	420
Depleted.....	90	21	8	720
Total.....	129			2,240

There is no dense growth within the meaning of the definition adopted in this report—that is, bottom on which 8 or more bushels of market oysters may be tonged by a man working 9 hours. There are two areas of scattering growth which lie as strips along the line of a shoal largely exposed at low water. These areas are sufficiently productive to yield from 5 to 10 bushels of market oysters per day to the tonger, the average being about 7 bushels.

The only other productive bottom is a very scattering area occupying the central and eastern portion of the rock, from the ridge to the deep water lying between this shoal and High Shoal. On this area a tonger can average a little over 3 bushels of oysters per day. Depleted areas lie on each side of the shoal, that on the western side being more extensive and continuous with the depleted area of Dog Shoal rock. On these areas a tonger could take hardly a bushel of market oysters per day, although there are spots a little more productive. On the scattering and very scattering areas of market oysters the young growth is prolific, on the former being sufficient to yield to the tonger an average of about 26 bushels per day and on the latter about 16 bushels. On both of these areas there is an abundance of clean shells suitable for taking a set of spat, but the depleted areas have comparatively few shells and young oysters.

The following observations were made on this rock:

DETAILS OF EXAMINATION OF TROUT SHOAL ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
		<i>Fect.</i>					<i>Bush.</i>	<i>Bush.</i>	<i>Bush.</i>
76	Aug. 13, 1909	6.5	Scattering.....	7.7	32.3	4.0	260	48	308
105	Aug. 14, 1909	4.0	do.....	1.0	16.1	2.4	111	29	140
482	Sept. 8, 1909	4.5	do.....	4.7	29.1	4.9	220	59	279
488	do.....	6.5	do.....	4.2	21.6	3.4	168	41	209
75	Aug. 13, 1909	6.0	Very scattering....	4.3	18.3	2.0	147	24	171
106	Aug. 14, 1909	5.0	do.....	.8	25.6	1.8	171	22	193
107	do.....	10.0	do.....	.4	18.7	3.3	124	40	164
108	do.....	9.5	do.....	.0	4.8	2.8	31	34	65
72	Aug. 13, 1909	11.5	Depleted.....	.4	2.1	1.7	16	21	37
483	Sept. 8, 1909	11.0	do.....	.7	2.3	.3	19	4	23
484	do.....	5.5	do.....	1.6	6.1	.6	50	7	57
487	do.....	6.5	do.....	.0	.0	.0	0	0	0

DOG SHOAL ROCK.

This bed occupies the northwestern or upstream part of Naseway Shoal. The rocks are in reality two, separated by the tongue of deeper water which makes into Naseway Shoal from the west and extends well toward the ridge of Trout Shoal.

The larger rock is hook shaped and contains two areas of dense growth and a long strip of very scattering oysters, both following the line of a shell ridge bare in parts at low water. The smaller area is a U-shaped ridge of scattering oysters lying between the deeper water just mentioned and the swash channel, which separates it from the adjacent Fishing Point rock.

The following table shows the area, density of growth, and estimated oyster content of the rock:

OYSTER GROWTH ON DOG SHOAL ROCK.

Character of growth of market oysters.	Area.	Oysters per acre.		Estimated content of market oysters.
		Seed.	Market.	
	<i>A cres.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	16	155	104	1,664
Scattering.....	13	153	39	507
Very scattering.....	35	22	27	945
Depleted.....	118	41	12	1,416
Total.....	182			4,532

The dense areas produce market oysters in sufficient quantity to yield the tonger an average of about 12 bushels per day, the scattering area will yield about 6 bushels, and the very scattering about 3 bushels. On the depleted area the yield would be at no place more than about 2 or 2½ bushels of marketable stock, and the average at all places examined was about 1 bushel.

The growth of young oysters on this rock is prolific, the density on the dense and scattering areas of market oysters being sufficient to yield the tonger an average of about 23 bushels per day. On the area of very scattering growth the yield should be about 3 bushels of young per day and on the depleted area about 4 bushels. The average of the latter is brought up by the very dense growth of young found in places close to the exposed ridge, where the quantity of market oysters was negligible. Over all of the area shown on the chart as depleted, excepting close to the productive areas, both clean shells and young were practically absent. The market oysters on this rock, like those on Trout Shoal and High Shoal, are comparatively small, averaging a little in excess of 400 per bushel.

The data on which the foregoing description is based are as follows:

DETAILS OF EXAMINATION OF DOG SHOAL ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
		<i>Fect.</i>					<i>Bush.</i>	<i>Bush.</i>	<i>Bush.</i>
79	Aug. 13, 1909	7.0	Dense	10.2	30.3	9.1	263	110	373
103	Aug. 14, 1909	5.5	do	3.7	21.0	6.0	160	73	233
104	do	5.5	do	4.2	21.2	3.0	165	36	201
491	Sept. 8, 1909	6.0	do	7.4	21.3	16.0	187	194	381
493	do	9.5	do	.0	.0	9.0	0	109	109
102	Aug. 14, 1909	4.5	Scattering	1.2	15.2	2.4	106	29	135
465	Sept. 3, 1909	6.0	do	9.2	21.7	4.1	201	49	250
78	Aug. 13, 1909	4.0	Very scattering	2.4	3.2	1.0	36	12	48
486	Sept. 8, 1909	8.0	do	.0	4.5	2.7	29	33	62
492	do	8.5	do	.0	2.0	3.0	13	36	49
494	do	7.0	do	.0	.0	2.1	0	25	25
496	do	6.5	do	.6	3.0	2.4	23	29	53
77	Aug. 13, 1909	7.5	Depleted	4.1	11.0	1.6	98	19	117
82	do	5.5	do	6.4	42.1	1.3	302	16	318
83	do	9.0	do	.3	.0	2.3	2	28	30
101	Aug. 14, 1909	10.0	do	.0	.3	.7	2	8	10
485	Sept. 8, 1909	11.5	do	.0	.3	2.3	2	28	30
487	do	6.5	do	.0	.0	.0	0	0	0
490	do	8.5	do	.0	.0	1.7	0	21	21
495	do	7.0	do	.0	.3	.0	2	0	2
500	do	6.5	do	.0	.3	.0	2	0	2
501	do	6.5	do	.0	.0	.0	0	0	0

FISHING POINT ROCKS.

These are two beds of considerable extent lying between High Shoal and Naseway Shoal and Fishing Point. The names by which they are known to the oystermen were not learned. One of these beds, embracing scattered and very scattered areas, lies between High Shoal and the Baylor line, almost surrounded by deeper water; the other, which includes dense, scattered, and depleted areas, lies along the Baylor line inside of Naseway Shoal, from which it is separated by a channel carrying from 8 to 10 feet at low water.

The statistics of the rocks are as follows:

OYSTER GROWTH ON FISHING POINT ROCKS.

Character of growth of market oysters.	Area.	Oysters per acre.		Estimated content of market oysters.
		Seed.	Market.	
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense	45	185	119	5,355
Scattering	77	178	82	6,314
Very scattering	47	70	44	2,068
Depleted	90	30	19	1,710
Total	259			15,447

It is estimated that on the dense area a tonger could take an average of 10 or 11 bushels of market oysters per day, and on the scattering area about 7 bushels. On the very scattering area the water is rather deep, the beds in this vicinity ranging from about 12 to 22 feet at low water, and although the density of growth is fair as compared with other beds described, this reduces the probable yield to the tonger to an average of between 3 and 4 bushels per day. On the depleted

area the probable average yield is estimated at between $1\frac{1}{2}$ and 2 bushels per day of tonging. On the area of scattering growth on the bed inside of High Shoal young growth is almost absent, giving the bottom the appearance of having been planted. On the very scattering area in the same bed the quantity of young is sufficient to yield about 6 bushels per day per tonger.

On the dense and scattering areas of the other bed there is a dense growth of young oysters, sufficient to yield on the former about 18 bushels and on the latter about 13 bushels per day. On the depleted bottom as a whole it would probably be impossible to take more than 3 bushels of young per day, though there are spots where the yield might be double that amount.

The following results were obtained from examinations:

DETAILS OF EXAMINATION OF FISHING POINT ROCKS.

Station number.	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
		<i>Feet.</i>					<i>Bush.</i>	<i>Bush.</i>	<i>Bush.</i>
466	Sept. 3, 1909	8.0	Dense.....	3.8	27.4	8.5	203	103	306
467do.....	8.0do.....	1.2	24.7	11.2	168	135	303
62	Aug. 13, 1909	14.0	Scattering.....	.0	.3	6.7	2	81	83
468	Sept. 3, 1909	9.0do.....	6.2	39.7	6.1	298	74	372
469do.....	9.5do.....	2.5	18.3	7.5	135	91	226
66	Aug. 13, 1909	13.0	Very scattering.....	4.0	6.8	3.6	70	44	114
73do.....	8.5	Depleted.....	2.6	6.9	2.0	62	24	86
84do.....	10.0do.....	.0	1.3	.7	8	8	16
99	Aug. 14, 1909	9.0do.....	.3	7.3	2.4	49	29	78
100do.....	8.0do.....	.3	.3	1.2	4	14	18

ROCKS BETWEEN FISHING POINT AND BALLARDS MARSH ROCKS.

In this region there are two small rocks for which no names were obtained. One of these lies close to the Baylor line and is encroached on by planted areas. It consists of a dense area inshore, the oysters becoming very scattering farther out, surrounded by a fringe of depleted bottom. The other bed is a small patch of very scattering growth about 400 yards farther out, in the direction of Dog Shoal Rock.

The following table exhibits the extent and condition of these beds:

OYSTER GROWTH ON ROCKS BETWEEN FISHING POINT AND BALLARDS MARSH ROCKS.

Character of growth of market oysters.	Area.	Oysters per acre.		Estimated content of market oysters.
		Seed.	Market.	
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	5	183	268	1,340
Very scattering.....	8	1	31	248
Depleted.....	18	0	0	0
Total.....	31			1,588

The dense area varies considerably in productiveness, one spot near what appeared to be the center of the original bed producing a quantity of oysters sufficient to yield 50 bushels per day to the tonger, while in another place not more than 9 bushels could be taken.

On the very scattering area of the larger bed barely 3 bushels per day could be taken, but on the small isolated spot the growth was sufficient to yield about 4 bushels per day. The depleted area is practically bare of oysters of all sizes, and the quantity of shells is negligible. On the small strip in the dense bottom where the market oysters were most abundant there is a very dense growth of young, but the rest of the bed is deficient in this respect.

The following table gives the results of the several examinations of the beds:

DETAILS OF EXAMINATION OF BEDS BETWEEN FISHING POINT AND BALLARDS MARSH ROCKS.

Station number.	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
		<i>Feet.</i>					<i>Bush.</i>	<i>Bush.</i>	<i>Bush.</i>
93	Aug. 14, 1909	7.5	Dense	0.0	0.0	7.0	0	85	85
95do.....	7.5do.....	.0	56.4	37.3	366	451	817
98do.....	7.0	Very scattering.....	.3	.0	3.1	2	37	39
464	Sept. 3, 1909	8.5do.....	.0	.0	2.0	0	25	25
96	Aug. 14, 1909	8.5	Depleted.....	.0	.0	.0	0	0	0
97do.....	8.5do.....	.0	.0	.0	0	0	0
463	Sept. 3, 1909	8.5do.....	.0	.0	.0	0	0	0

BALLARDS MARSH ROCK.

This is the bed called by Winslow "Bally Smash," probably an unconscious attempt to render a provincial pronunciation phonetically. It is the westernmost bed of Public Ground No. 6, Isle of Wight County. It follows the line of a shoal which sets offshore from Ballards Marsh.

Its extent and condition are epitomized in the following table:

OYSTER GROWTH ON BALLARDS MARSH ROCK.

Character of growth of market oysters.	Area.	Oysters per acre.		Estimated content of market oysters.
		Seed.	Market.	
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Scattering.....	4	152	31	124
Very scattering.....	35	191	24	792
Depleted.....	142	45	7	894
Total.....	179			1,810

The scattering area is a small spot near the inner end of the shoal, where the quantity of market oysters is sufficient to yield to the tonger between 7 and 8 bushels per day. The very scattering growth is found on each side of this and beyond it for a distance of about two-thirds the length of the shoal, bearing a growth yielding about 3½ bushels of oysters per day's work. On both sides of the very scattering area and beyond it along the line of the shoal is a depleted bottom on which, as a whole, less than 1 bushel of oysters can be taken per day, the edges of the area being practically barren.

Along both sides of the shoal, even on the so-called depleted bottom which surrounds it at its outer end, is a heavy growth of young and many shells, which will yield on the average about 35 bushels of culls and spat per day. The market oysters on this bed will average about 400 to the bushel and the culls or seed oysters about 750.

DETAILS OF EXAMINATION OF BALLARDS MARSH ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
		<i>Feet.</i>					<i>Bush.</i>	<i>Bush.</i>	<i>Bush.</i>
86	Aug. 14, 1909	3.0	Scattering	13.7	9.7	2.6	152	31	183
85	do	6.0	Very scattering	3.2	3.2	2.1	42	25	67
462	Sept. 3, 1909	5.0	do	6.6	21.7	2.0	184	25	209
514	Sept. 9, 1909	4.0	do	10.8	42.3	1.8	346	22	368
87	Aug. 14, 1909	6.0	Depleted3	2.3	1.4	17	17	34
88	do	7.0	do0	.0	.0	0	0	0
89	do	9.0	do0	.0	1.7	0	21	21
90	do	7.0	do0	.0	.0	0	0	0
91	do	7.0	do0	.3	1.8	2	22	24
461	Sept. 3, 1909	7.0	do	4.7	27.7	1.3	211	16	227
504	Sept. 9, 1909	6.0	do0	.0	.0	0	0	0
505	do	5.0	do	5.7	22.5	.4	183	5	188
506	do	4.5	do	7.4	21.0	.5	185	6	191
508	do	4.5	do	10.3	22.4	.2	213	2	215
513	do	6.0	do0	.0	.2	0	2	2
515	do	5.5	do0	.0	.9	0	11	11
516	do	6.0	do0	.0	.7	0	8	8
517	do	6.0	do0	.0	.2	0	2	2
518	do	7.0	do0	.0	.5	0	6	6
519	do	7.0	do0	.3	1.4	2	17	19
520	do	6.5	do0	.0	.0	0	0	0
521	do	5.5	do0	.0	.0	0	0	0

CREEK CHANNEL ROCK.

This is a small bed about 2 acres in extent, covering a shoal marked by a buoy of the Light-House Establishment. It has the indications of having been a dense bed, but at the present time it is depleted, and a tonger could take on it an average of hardly a bushel of oysters per day, and the young growth is still more sparse. It is surrounded by private beds. It constitutes Public Ground No. 5, Isle of Wight County.

The following observations were made on this bed:

DETAILS OF EXAMINATION OF CREEK CHANNEL SHOAL ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
383	Aug. 26, 1909	<i>Feet.</i> 13.0	Depleted.....	0.0	0.0	1.4	<i>Bushels.</i> 0	<i>Bushels.</i> 19	<i>Bushels.</i> 19
384do.....	6.0do.....	0.0	1.1	.2	7	3	10

AARON SHOAL ROCK.

This is the only bed in Public Ground No. 2, Isle of Wight County. It is almost or quite surrounded by private beds, the boundary stakes of which formed a forest which made it difficult to tell, without spending on the bed more time than its importance warranted, what was planted ground and what was not.

The following statistics exhibit its present condition:

OYSTER GROWTH ON AARON SHOAL ROCK.

Character of growth of market oysters.	Area.	Oysters per acre.		Estimated content of market oysters.
		Seed.	Market.	
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	2?	129	100	200
Scattering.....	4	135	33	132
Very scattering.....	3	0	23	69
Depleted.....	24	112	5	120
Total.....	31			521

The dense area forms a very narrow strip along the northern edge close to and among the stakes. Its area could not be very definitely determined without wasteful expenditure of time, but is probably about 2 acres. About 10 bushels of oysters per day could be taken by the tonger.

On the scattering growth it is estimated that about 4½ bushels, and on the very scattering about 3 bushels, per day could be taken. The depleted area is for the most part bare. There is a good growth of young on the dense and scattered areas and at two spots on the depleted bottom.

The following examinations were made.

DETAILS OF EXAMINATION OF AARON SHOAL ROCK

Station number.	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
		<i>Feet.</i>					<i>Bush.</i>	<i>Bush.</i>	<i>Bush.</i>
392	Aug. 26, 1909	8.5	Dense.....	0.0	6.0	9.7	39	134	173
396	do.....	5.5	do.....	5.7	28.0	4.7	219	65	284
389	do.....	5.5	Scattering.....	1.6	19.2	2.4	135	33	168
400	do.....	6.0	Very scattering.....	.0	.0	1.7	0	23	23
390	do.....	5.0	Depleted.....	6.7	24.2	1.3	201	18	219
391	do.....	8.5	do.....	5.0	10.3	.3	90	4	94
394	do.....	7.5	do.....	.0	.0	.0	0	0	0
395	do.....	6.5	do.....	5.3	18.7	.0	156	0	156

BROWNS SHOAL ROCKS.

Included under this name are a number of small rocks, separated by depleted and barren bottom lying at the extreme lower end of public ground No. 1, Warwick County, just above Newport News. The productive portions lie on Browns Shoal and a number of other shoal spots in the vicinity. The extent and present condition of the rocks as a whole are shown in the following table:

OYSTER GROWTH ON BROWNS SHOAL ROCKS.

Character of growth of market oysters.	Area.	Oysters per acre.		Estimated content of market oysters.
		Seed.	Market.	
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	68	126	183	12,444
Scattering.....	44	142	54	2,376
Very scattering.....	27	88	39	1,053
Depleted.....	226	5	4	904
Total.....	365			16,777

The dense area is found in seven patches, of which the largest, covering about 25 acres, is on a shoal west of Browns Shoal surrounding a watchhouse or covered pierhead. The areas as a whole are quite productive of market oysters, and it is estimated that an industrious tonger working nine hours per day could take between 10 and 40 bushels of oysters, the average at all places examined being about 15 bushels.

The areas of scattering growth are three in number, lying on the ends of Browns Shoal proper and a small shoal west of it, inshore of the watchhouse above alluded to. They carry a depth of between about 4 or 5 and 12 feet at low water, and their productiveness is such that a tonger could take an average of between 5 and 6 bushels of market oysters per day.

The areas of very scattering growth are a number of small patches nearly all lying between the more prolific areas and the edges of the bed. They are nearly all covered by about 10 feet of water at low tide, and bear oysters in sufficient quantity to yield to the tonger between 3 and 4 bushels per day.

The depleted bottom constitutes nearly two-thirds of the total area of the beds as charted. At no place does it promise to yield during the present season more than 2½ bushels per day, and the major part of it is practically barren.

The growth of young oysters is very good on the shallower parts of the beds, especially on those portions yielding a scattering growth of market stock, where a tonger could take an average of about 15 bushels per day. On the dense areas as a whole the young growth is less abundant, the estimated average yield being about 11 bushels per day, the heaviest growth being on two small shoals between the inner end of Browns Shoal and the shipyard at Newport News.

On all of the productive areas there is an abundance of shells suitable for catching a set of spat, but the depleted areas are practically bare and give no promise of recuperation under any natural conditions.

The following exhibits the results of examinations:

DETAILS OF EXAMINATION OF BROWNS SHOAL ROCKS.

Station number.	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
		<i>Feet.</i>					<i>Bush.</i>	<i>Bush.</i>	<i>Bush.</i>
112	Aug. 16, 1909	9.5	Dense	4.7	31.9	12.0	238	165	403
122do.....	9.5do.....	.0	24.6	7.9	160	109	269
123do.....	8.5do.....	1.3	21.6	9.6	149	133	282
126do.....	8.0do.....	.0	10.7	7.3	70	101	171
139	Aug. 17, 1909	9.5do.....	1.6	6.4	32.0	52	443	495
141do.....	11.5do.....	2.2	11.2	9.6	86	133	219
142do.....	14.5do.....	7.2	15.6	12.8	148	177	325
144do.....	10.5do.....	1.5	18.0	10.5	127	145	272
145do.....	12.5do.....	1.5	11.0	14.5	81	200	281
146do.....	13.5do.....	.0	9.2	24.4	60	337	397
450	Sept. 1, 1909	10.0do.....	5.2	37.8	4.4	280	61	341
459	Sept. 3, 1909	10.5do.....	5.0	21.2	10.8	170	149	319
460do.....	10.5do.....	2.1	16.2	20.8	119	287	406
524	Sept. 10, 1909	11.5do.....	.0	4.0	9.0	26	124	150
115	Aug. 16, 1909	9.0	Scattering	.3	28.7	6.3	188	87	275
118do.....	7.0do.....	.6	21.2	4.0	141	55	196
119do.....	6.0do.....	.9	12.8	2.3	89	32	121
120do.....	6.0do.....	4.5	11.8	2.5	100	34	134
136	Aug. 17, 1909	10.0do.....	3.6	25.6	4.4	190	61	251
138do.....	9.0do.....	2.0	31.0	4.0	214	55	269
143do.....	12.5do.....	.0	2.8	6.4	18	88	106
451	Sept. 1, 1909	7.5do.....	12.3	34.0	3.0	301	41	342
452do.....	11.0do.....	7.3	20.4	5.8	180	80	260
453do.....	10.0do.....	2.9	12.1	4.6	98	64	162
117	Aug. 16, 1909	5.0	Very scattering	.2	12.9	2.2	85	30	115
125do.....	10.0do.....	.3	3.7	3.3	26	45	71
135	Aug. 17, 1909	11.0do.....	2.0	6.0	3.2	52	44	96
140do.....	11.0do.....	0	8.8	2.8	57	39	96
457	Sept. 3, 1909	11.0do.....	8.2	26.2	2.5	223	35	258
458do.....	10.5do.....	2.9	9.6	3.8	81	52	133
523	Sept. 10, 1909	10.0do.....	2.0	12.5	2.0	94	28	122
113	Aug. 16, 1909	12.6	Depleted	.0	5.7	2.0	37	28	65
114do.....	10.5do.....	.0	.0	.0	0	0	0
116do.....	12.0do.....	.0	.3	.0	2	0	2
124do.....	12.0do.....	.0	.0	.0	0	0	0
137	Aug. 17, 1909	12.0do.....	.0	.0	.0	0	0	0
449	Sept. 1, 1909	18.0do.....	.0	.0	.0	0	0	0
454do.....	18.0do.....	.0	2.3	1.4	15	19	34
455do.....	10.5do.....	.0	.0	.0	0	0	0
456do.....	11.0do.....	.0	.0	.0	0	0	0
522	Sept. 10, 1909	24.0do.....	.0	.0	.0	0	0	0
567	Sept. 14, 1909	20.5do.....	.0	.0	.0	0	0	0

GUN ROCK.

This is a small bed lying on a shoal spot west of the preceding. Its extent and estimated density of growth and contents are as follows:

OYSTER GROWTH ON GUN ROCK.

Character of growth of market oysters.	Area.	Oysters per acre.		Estimated content of market oysters.
		Seed.	Market.	
	Acres.	Bushels.	Bushels.	Bushels.
Dense.....	6	198	152	912
Very scattering.....	16	62	30	480
Depleted.....	4	0	0	0
Total.....	26			1,392

The dense area forms a narrow tongue running along the inner or shoreward end of the ridge or backbone of the shoal, and it produces market oysters in sufficient quantity to yield the tonger about 9 bushels per day. There is no scattering growth, but the outer half of the length of the bed as far as the Baylor line produces a very scattering growth sufficient to yield about 3 bushels per day.

Along the higher parts of the ridge, on both the dense and very scattering bottoms, there is a prolific growth of young oysters, sufficient to furnish the tonger from 12 to 14 bushels per day. The edges of the very scattering area produce but few young oysters, and the depleted bottom is practically bare of both oysters and clean shells, its position being indicated solely by the presence of shells more or less deeply buried in the mud.

The following observations were made:

DETAILS OF EXAMINATION OF GUN ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
		<i>Feet.</i>					<i>Bush.</i>	<i>Bush.</i>	<i>Bush.</i>
134	Aug. 17, 1909	16.0	Dense.....	2.0	28.5	11.0	198	152	350
147	do.....	6.5	Very scattering.....	.8	17.4	1.8	118	25	143
149	do.....	12.5	do.....	.0	1.0	2.5	6	34	40
148	do.....	12.5	Depleted.....	.0	.0	.0	0	0	0

KETTLE HOLE ROCK.

I am not certain of the name of this bed, as in the field some confusion arose as to whether this or the next was Thomas Point Rock. In designating it as above I have been guided by Winslow's nomenclature. This bed is the largest and most important in this part of

the river. It lies on the north side of a shallow ridge, extending thence shoreward toward Watts Creek as far as the Baylor line.

Its extent and present general condition are as follows:

OYSTER GROWTH ON KETTLE HOLE ROCK.

Character of growth of market oysters.	Area.	Oysters per acre.		Estimated content of market oysters.
		Seed.	Market.	
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	258	317	105	27,090
Scattering.....	66	207	82	5,412
Very scattering.....	111	180	42	4,662
Depleted.....	11	0	0	0
Total.....	446			37,164

The principal area of dense growth starts from the ridge and extends as a broad belt as far as the Baylor line inshore. The public ground includes but a portion of the shoal, possibly because the southern part was taken up as private ground prior to the Baylor survey. The productive bottom probably extends across the ridge, but as it is not included in the public grounds and is staked as a private holding it was not examined. The growth of market oysters on the public ground is rather uniform and is sufficient to provide the tonger with from 8 to 13 bushels per day, the average being about 9 or 10. Near the middle of the inner part of the dense area is a small spot where market oysters are deficient in quantity, but the growth of young is so prolific that it has not been thought advisable to differentiate it on the chart. There is also a small area of dense growth on a shallow spot east of the main area.

The scattering growth on Kettle Hole Rock forms a fringe along the western border of the dense area, with a broad tongue thrust into the latter near its middle. The growth of market oysters is sufficient to yield to the tonger an average of about 6 bushels per day.

The areas of very scattering growth form a border on the eastern and inshore edges of the dense growth, and it is estimated that about 3 or 3½ bushels of oysters could be tonged per day on the areas taken in their entirety.

The depleted bottom is insignificant and bare of everything except a few buried shells.

There is a heavy growth of young oysters over practically the entire extent of this bed. On the dense areas they are estimated to be present in sufficient quantities to yield to the tonger about 28 bushels per day as an average at the beginning of the season, while on the scattering and very scattering areas the yield would probably average about 16 or 17 bushels. Undoubtedly the entire bed can be regarded as

presently and prospectively productive. The following data were obtained from the examinations made:

DETAILS OF EXAMINATION OF KETTLE HOLE ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
		<i>Feet.</i>					<i>Bush.</i>	<i>Bush.</i>	<i>Bush.</i>
133	Aug. 17, 1909	10.0	Dense	8.5	47.5	10.0	364	138	502
155	do	13.5	do	5.0	62.0	7.5	436	104	540
156	do	11.5	do	1.5	29.9	7.8	204	108	312
157	do	11.5	do0	32.0	8.0	208	111	319
159	do	9.5	do0	44.6	.7	290	10	300
360	Aug. 26, 1909	11.5	do	5.0	27.5	11.7	211	161	372
361	do	11.5	do	2.1	18.5	7.6	134	105	239
362	do	10.5	do	8.1	54.6	9.6	408	133	541
363	do	8.5	do	6.5	47.2	6.1	349	84	433
364	do	7.5	do	7.6	46.7	7.9	353	109	462
365	do	8.0	do	9.6	72.4	6.5	533	90	623
158	Aug. 17, 1909	12.5	Scattering	5.0	34.6	5.5	257	76	333
163	Aug. 18, 1909	17.0	do7	50.0	6.9	334	95	429
359	Aug. 26, 1909	11.0	do9	3.8	5.4	31	75	106
151	Aug. 17, 1909	11.0	Very scattering0	16.0	4.0	104	55	159
152	do	13.0	do4	43.3	2.1	284	29	313
153	do	11.5	do	1.1	13.3	3.0	95	41	136
154	do	11.5	do2	24.0	3.3	169	46	215
369	Aug. 26, 1909	11.5	do	7.9	30.0	2.9	246	40	286
164	Aug. 18, 1909	14.0	Depleted0	.0	.0	0	0	0
165	do	16.0	do0	.0	.0	0	0	0

THOMAS POINT ROCK.

As explained in the discussion of the preceding bed, there may be some question as to the name of this one, which lies between what has been called Kettle Hole Rock and Blunt Point Rock. It is entirely separated from the former by a swash channel carrying from 12 to 21 feet of water, but is connected with Blunt Point Rock by a narrow ridge of depleted bottom. The bed lies on and about two shoals which extend from the edge of the channel lying north of White Shoal Light. Its extent and condition are as follows:

OYSTER GROWTH ON THOMAS POINT ROCK.

Character of growth of market oysters.	Area.	Oysters per acre.		Estimated content of market oysters.
		Seed.	Market.	
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense	76	168	115	8,745
Scattering	118	170	71	8,378
Very scattering	100	103	51	5,100
Depleted	127	80	21	2,667
Total	421			24,890

The dense growth of market oysters is in three patches, all lying on or close to the shoaler parts of the bed in water ranging from 5 to 8 feet at low tide. The growth is sufficiently prolific to yield to the tonger about 10 bushels of marketable oysters per day.

The area covered by scattering marketable oysters stretches from the western side of the longer shoal across some intervening deeper water to two small shoals to the westward. Over the whole area the density of growth is such that about 6 bushels of oysters may be taken per day.

The very scattered growth is in three areas fringing the denser parts of the bed. Its productiveness varies between areas which will yield $2\frac{1}{2}$ and those which will yield $4\frac{1}{2}$ bushels per day, the general average at all places examined being about $3\frac{1}{2}$ bushels.

The best of the depleted bottom, which is in the areas lying on the edge of the deep-water channel, will yield about $2\frac{1}{2}$ bushels per day, while the inshore area and that lying in the midst of the scattering growth will not yield an average of over 1 bushel.

The young growth is in good quantity, though not so abundant as on Kettle Hole Rock. On the dense and scattering areas it is sufficient to yield an average of about 15 bushels per day. On the very scattering area near the inshore end of the eastern edge of the rock it is in about the same abundance, but elsewhere it will yield not more than $4\frac{1}{2}$ bushels per day's tonging. On the best of the depleted bottom, along the edge of the deep-water channel, it is estimated that about 10 or 11 bushels per day could be taken by the tonger, but on other parts of the depleted area young oysters are practically absent.

The oysters on this bed, as on all others on this shore of the river, are of fair size, the marketable stock averaging about 350 per bushel and the young approximately 750.

The following observations were made:

DETAILS OF EXAMINATION OF THOMAS POINT ROCK.

Station number	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
		<i>Feet.</i>					<i>Bush.</i>	<i>Bush.</i>	<i>Bush.</i>
127	Aug. 17, 1909	12.5	Dense.....	0.8	2.1	11.2	19	154	173
132do.....	12.5do.....	9.6	33.0	8.9	277	123	400
366	Aug. 26, 1909	8.5do.....	3.9	33.1	10.3	241	142	383
367do.....	8.5do.....	13.4	13.8	5.8	177	80	257
374do.....	13.0do.....	5.8	13.6	5.5	126	76	202
128	Aug. 17, 1909	14.5	Scattering.....	2.0	21.2	6.0	151	83	234
129do.....	12.5do.....	2.5	24.0	4.5	172	62	234
130do.....	13.5do.....	1.5	9.5	5.5	71	76	147
131do.....	9.5do.....	5.5	24.0	5.5	192	76	268
160	Aug. 18, 1909	9.0do.....	3.0	31.1	5.7	221	79	300
161do.....	9.0do.....	2.9	27.7	5.1	199	70	269
368	Aug. 26, 1909	8.5do.....	10.0	28.6	1.8	251	25	276
372do.....	11.0do.....	2.3	13.2	7.1	101	98	199
162	Aug. 18, 1909	13.0	Very scattering.....	1.2	13.5	2.4	97	38	130
166do.....	15.5do.....	2.7	45.9	5.5	316	76	392
371	Aug. 26, 1909	14.0do.....	.0	4.5	3.3	29	46	75
375do.....	17.0do.....	.0	6.8	3.6	44	50	94
376do.....	13.0do.....	.0	4.5	3.8	29	52	81
168	Aug. 18, 1909	9.5	Depleted.....	1.3	18.5	1.8	129	25	154
172do.....	15.0do.....	2.7	24.6	2.7	177	37	214
373	Aug. 26, 1909	13.0do.....	.0	1.2	.9	8	12	20
377do.....	13.0do.....	.3	1.0	.7	8	10	18

BLUNT POINT ROCK.

This is the uppermost bed in that part of Public Ground No. 1, Warwick County, in which the cull law is enforced. All beds above this, excepting the small one in Warwick River, are within the area which is set apart for seed production. This rock is rather attenuated in most of its parts, being in the shape of an irregular ring surrounding a deeper barren area, with a long tail running along the Baylor line in the direction of Deep Creek. The highly productive area is very limited and the very scattering growth constitutes more than half of the total area. The extent and general condition of the bed at the time of the survey are shown in the following table:

OYSTER GROWTH ON BLUNT POINT ROCK.

Character of growth of market oysters.	Area.	Oysters per acre.		Estimated content of market oysters.
		Seed.	Market.	
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	16	171	161	2,576
Scattering.....	69	193	60	4,140
Very scattering.....	225	105	45	10,125
Depleted.....	118	42	14	1,652
Total.....	428			18,493

The bottom covered with a dense growth occurs on two small patches on small shoals, on which about 15 bushels of market oysters could be taken in a day.

There are three areas of scattering growth, of about equal productiveness so far as market oysters are concerned. One of these is at the extreme end of the rock off Deep Creek, which is in close proximity to planted beds and bears some indications of being itself planted ground. These areas as a whole will yield, it is estimated, an average of about 6 bushels per day.

The very scattering growth consists of a narrow zone almost encircling the included barren area above alluded to and a prolongation northward toward Deep Creek. The examinations made on it indicate a probable yield of about $3\frac{1}{2}$ bushels per day of continuous tonging.

The depleted area skirts the preceding for a good part of its length, and in addition forms a projection on the western part of the bed and a small isolated patch on a shoal just beyond it. Its content of oysters is such that it could furnish the tonger with hardly more than a bushel per day.

The young growth is in good quantity on the dense and scattering areas of oysters, excepting that nearest Deep Creek, where it is practically absent. With the exception noted, the tonger should be able to gather about 15 bushels per day. On the narrow annular part of

the beds the production of young oysters is good on the very scattering and depleted areas, which in their other parts are deficient in immature growth.

The following data furnish the basis for the foregoing:

DETAILS OF EXAMINATION OF BLUNT POINT ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
		<i>Fect.</i>					<i>Bush.</i>	<i>Bush.</i>	<i>Bush.</i>
252	Aug. 21, 1909	9.5	Dense.....	7.9	17.5	7.5	165	103	268
253do.....	12.5do.....	3.1	24.5	15.9	178	219	397
180	Aug. 18, 1909	12.5	Scattering.....	1.5	33.9	5.3	230	73	303
187do.....	6.5do.....	.0	.8	3.5	5	48	53
254	Aug. 21, 1909	11.0do.....	.3	22.9	4.2	151	58	209
173	Aug. 18, 1909	16.0	Very scattering....	.9	5.0	4.6	38	63	101
175do.....	10.5do.....	2.6	42.3	3.5	292	48	340
178do.....	10.5do.....	1.3	27.1	2.9	184	40	224
182do.....	11.0do.....	2.0	7.1	3.7	59	51	110
188do.....	9.5do.....	.0	4.0	2.0	26	28	54
380	Aug. 26, 1909	14.5do.....	1.0	3.1	4.1	27	57	84
381do.....	13.0do.....	.3	16.0	2.1	106	29	135
174	Aug. 18, 1909	16.0	Depleted.....	1.9	35.2	2.6	241	36	277
177do.....	13.5do.....	.0	.0	.0	0	0	0
179do.....	16.5do.....	.0	3.6	.9	23	7	30
181do.....	13.5do.....	.0	.0	.0	0	0	0
189do.....	8.5do.....	1.3	12.1	1.3	87	18	105
190	Aug. 19, 1909	9.5do.....	.0	.0	1.3	0	18	18
251	Aug. 21, 1909	11.5do.....	.8	2.9	1.3	24	18	42
378	Aug. 26, 1909	12.5do.....	.0	.0	1.8	0	25	25
379do.....	15.5do.....	.0	.0	.0	0	0	0

WHITE SHOAL ROCKS.

These are two rocks in very shallow water, with slightly greater depths between and deep channels surrounding. The westernmost lies about a bare shell bank and the easternmost is nearly awash at low water.

The following exhibits their extent and condition at the beginning of the oyster season of 1909-10.

OYSTER GROWTH ON WHITE SHOAL ROCKS.

Character of growth of market oysters.	Area.	Oysters per acre.		Estimated content of market oysters.
		Seed.	Market.	
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	44	312	127	5,588
Very scattering.....	10	108	36	360
Depleted.....	52	53	12	624
Total.....	106			6,572

The dense areas produce a good quantity of marketable oysters and at the beginning of the present season should be capable of yielding about 12 bushels of oysters per full day of tonging. There are no areas of scattered growth within the definition of this report, but at

each end of the western rock there is a small patch of very scattering growth capable of yielding an average of $3\frac{1}{2}$ bushels of marketable oysters per day.

The area charted as depleted bears very few marketable oysters.

The young growth on the dense areas is very prolific, being in sufficient quantity to afford a daily return to the tonger of about 35 bushels. On the scattering bottom about 10 bushels per day could be taken, while on the depleted bottom as a whole the average would hardly exceed 4 bushels, although two or three times that many could be taken in places.

DETAILS OF EXAMINATION OF WHITE SHOAL ROCKS.

Station number.	Date of examination.	Mean depth of water.	Character of growth of market oysters.	Oysters caught per square yard.			Estimated quantity oysters per acre.		
				Spat.	Culls.	Counts.	Seed.	Market.	Total.
		<i>Fect.</i>					<i>Bush.</i>	<i>Bush.</i>	<i>Bush.</i>
169	Aug. 18, 1909	5.5	Dense	7.7	92.3	6.5	650	90	740
170	do	8.5	do	3.8	10.4	9.2	92	127	219
443	Sept. 1, 1909	10.0	do	11.7	63.3	7.1	487	98	585
444	do	10.0	do	11.7	29.6	17.9	268	247	515
445	do	13.0	do	15.2	26.9	8.3	274	115	389
447	do	9.5	do	10.0	28.4	7.1	250	98	348
556	Sept. 14, 1909	9.5	do	6.1	26.2	11.6	210	160	370
557	do	4.0	do	11.5	29.3	5.7	265	79	344
558	do	7.5	Very scattering	2.0	8.7	3.0	70	41	111
562	do	10.5	do	2.6	19.7	2.3	145	32	177
552	do	9.0	Depleted3	1.3	.3	10	4	14
553	do	8.5	do	1.0	6.3	1.0	47	14	61
554	do	6.5	do3	.0	.0	2	0	2
555	do	12.5	do	1.9	11.0	1.9	84	26	110
559	do	10.5	do	4.6	21.0	1.8	166	25	191
561	do	11.5	do3	.3	.3	4	4	8

SEED OYSTER AREA.

JAIL ISLAND ROCK.

For this and all of the following rocks the standard of density of growth is different from that adopted in the preceding descriptions. The cull law, so far as it relates to the size of oysters, does not apply, and oysters of whatever size may be taken. The entire content of the bed, both young and old, is therefore taken into consideration, and as the average price of seed oysters is about two-thirds of that of the market oysters from the James River, a larger quantity has to be taken to furnish a living wage. In all of the following descriptions a bed is regarded as dense when 12 or more bushels may be taken by a tonger in a day's work, as scattering when it will yield between 8 and 12 bushels, very scattering when it yields between 4 and 8, and depleted if less than 4 bushels can be tonged per day. As the market oysters sell for 45 cents and the seed oysters for but 30 cents per bushel, the financial return is essentially equal.

Jail Island Rock, which extends alongshore west of the mouth of Warwick River, is continuous at its offshore edge with Wreck Shoal

Rock, the boundary between them as adopted in this report being purely arbitrary.

The extent and general condition of the bed at the time of the survey was as follows:

OYSTER GROWTH ON JAIL ISLAND ROCK.

Character of growth.	Area.	Oysters per acre.	Estimated content of oysters.
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	227	143	32,461
Scattering.....	198	109	21,582
Very scattering.....	14	28	392
Depleted.....	508	8	4,064
Total.....	947	-----	58,499

The principal area of dense growth runs from the inner edge of Wreck Shoal Rock in a gradually narrowing belt to a tongue extending to within 200 or 300 yards of shore between Jail Island and the mouth of Warwick River. The depth of water gradually decreases from about 9 feet to 2 or 3 feet at low water, near Jail Island. There is also a small area of dense growth lying on an isolated patch in about 10 or 11 feet of water off the mouth of Warwick River, which, being just on the cull line, is arbitrarily included in the Jail Island bed for the purposes of this report. The dense bottom as a whole will afford the tonger an average catch per day of about 17 bushels of oysters of all sizes.

The area of scattering growth lies in a single body north and west of the preceding in from 6 to 11 feet of water. The growth on the whole is rather heavier in the deeper water, and as an average should yield approximately 9 bushels per day.

The very scattering growth is in a small patch immediately east of Jail Island, where the yield to the tonger should be about 5 bushels per day.

The depleted bottom forms a broad zone on the inshore side and a narrow strip on the eastern edge of the bed. There is also a depleted area adjoining the small, isolated, dense patch before described, and a small patch lying between that and the main bed. In most places the so-called "depleted bottom" is practically bare. There is but a moderate supply of shells on the dense area and on the scattering area close to it, but elsewhere the bed is deficient in this respect.

It is stated that the inshore portions of the bed, on the depleted bottom along the Baylor line, produce fine single oysters, which in calm weather are picked up one by one and bring a high price in the markets. The survey did not disclose any quantity of such oysters.

DETAILS OF EXAMINATION OF JAIL ISLAND ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.
				Spat.	Culls.	Counts.	
		<i>Feet.</i>					<i>Bushels.</i>
184	Aug. 18, 1909	11.0	Dense	1.3	49.0	4.6	306
198	Aug. 19, 1909	7.0	do	1.7	12.1	2.5	97
199	do	6.0	do0	8.5	6.9	138
200	do	4.0	do	1.5	8.8	4.2	110
404	Aug. 27, 1909	4.0	do	1.3	3.3	2.8	62
207	Aug. 19, 1909	10.0	Scattering	4.6	8.2	2.1	88
210	do	11.5	do	5.4	16.7	2.5	140
211	do	8.5	do0	2.7	6.2	99
402	Aug. 27, 1909	4.0	Very scattering	1.1	1.9	.9	28
183	Aug. 18, 1909	11.0	Depleted0	.0	.0	0
185	do	12.0	do4	1.7	1.3	32
192	Aug. 19, 1909	9.0	do0	.0	.7	8
208	do	7.5	do0	.0	.0	0
212	do	6.5	do0	.3	.3	6
401	Aug. 27, 1909	5.5	do7	.7	.0	7
403	do	4.0	do0	.0	.0	0

WRECK SHOAL ROCK.

This is a large, important, and productive bed extending from the preceding to the edge of deep water. For the purposes of this report, it is regarded as including the oyster growth on and about Wreck Shoal proper and the small shoal to the westward of its outer end. Excepting where it adjoins Jail Island bed, its boundaries are rather sharply defined by a sudden shoaling of the water. This is especially pronounced at the southern edge of the bed, where the bottom very abruptly rises from about 150 feet to within 6 feet of the surface. North of the smaller shoal the bed is prolonged into a narrow belt occupying a slightly shoaling ridge connected with a corner of Mulberry Swash Rock. The depth at low water varies from less than 5 feet on the shoals to 12 or 15 feet at the edges. On one small area projecting as a tongue from the southeast side the water reaches a maximum depth of 30 feet.

Wreck Shoal Rock is practically everywhere highly productive and no part of it falls below the standard here regarded as constituting denseness of growth. Accepting the arbitrary inner boundary here adopted, it has an area of about 506 acres. The oyster growth at the places examined ranges from 178 to 497 bushels per acre, the average being about 316. The heaviest growth is as a rule found on the shoaler places, which facilitates the removal of the product. This materially raises the average daily yield to the tonger, which ranges in different places from 12 bushels to 51 bushels, with a general average for the entire bed of over 29 bushels.

The bottom is well covered with clean shells and the bed can be regarded as being in a healthy and promising condition. In a few places there is a fair growth of large oysters and on the bed as a

whole the young growth over 1 inch in length is numerically more than double that under 1 inch.

The following table exhibits the data obtained from the several examinations made:

DETAILS OF EXAMINATION OF WRECK SHOAL ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.
				Spat.	Culls.	Counts.	
193	Aug. 19, 1909	<i>Fect.</i> 12.0	Dense	0.0	16.7	10.3	<i>Bushels.</i> 195
194do.....	9.0do.....	7.1	30.3	.0	191
195do.....	8.0do.....	2.5	24.6	3.8	178
196do.....	9.0do.....	23.7	34.6	.4	301
197do.....	6.5do.....	17.9	69.2	5.0	497
201do.....	12.0do.....	18.3	28.6	1.4	254
202do.....	11.0do.....	21.8	44.6	3.6	379
203do.....	11.0do.....	9.5	57.7	.0	343
204do.....	10.0do.....	26.7	46.9	.3	379
205do.....	7.0do.....	43.0	45.0	.4	456
206do.....	11.5do.....	15.8	18.9	4.2	222
209do.....	14.5do.....	12.7	31.8	2.3	252
428	Aug. 31, 1909	15.0do.....	3.6	30.0	2.7	200
429do.....	6.5do.....	38.1	41.2	.8	413
430do.....	8.5do.....	27.7	60.0	1.2	460
431do.....	16.5do.....	9.5	30.9	.4	210

DRY SHOALS ROCKS.

These are 5 small rocks lying west of the preceding on and about shoals which ebb nearly or quite bare. They are in general isolated and surrounded by deep water, though two of them are connected by narrow ridges of depleted bottom with Swash Rock and Mulberry Swash Rock, respectively.

Their present condition and extent are shown in the following table:

OYSTER GROWTH ON DRY SHOALS ROCKS.

Character of growth.	Area.	Oysters per acre.	Estimated total content of oysters.
Dense	<i>Acres.</i> 126	<i>Bushels.</i> 244	<i>Bushels.</i> 30,766
Scattering	18	124	2,232
Very scattering	9	85	765
Depleted	21	20	420
Total	174	34,183

Four of these rocks are composed wholly or in major part of bottom bearing a dense growth of oysters, while the fifth, the smallest, bears a scattering growth exclusively. The dense growth in its daily yield to the tonger varies, with the locality, between 13 and 59 bushels, the average density over the entire area being sufficient to permit a

daily catch of about 30 bushels per man. The average depth of water is about 5 to 7 feet at low tide.

The scattering growth is found in two places, one an isolated rock of small size and the other at the extremity of a larger bed where the productive bottom runs off to deeper water. There is not much difference in the density of the growth on the two places, and it is estimated that on the two a tonger could take an average of about 10 bushels of oysters per day.

The very scattering bottom lies in two small patches at opposite ends of the longest bed of the group, and the growth is so sparse as barely to remove the areas from the category of depleted bottom. The depleted areas are three in number, one in the deeper water at the tip of a rock and the others on the two connecting ridges mentioned earlier in this description. About the same numerical proportion exists between the culls and spat as on the preceding bed, and at one place on the dense area there is a good growth of marketable oysters averaging between 400 and 450 per bushel. There is a fair or good deposit of shells throughout the dense and scattering areas and on the apical area of very scattering growth, but elsewhere the rocks are deficient in this respect.

DETAILS OF EXAMINATION OF DRY SHOALS ROCKS.

Station number.	Date of examination.	Mean depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.
				Spat.	Culls.	Counts.	
		<i>Fect.</i>					<i>Bushels.</i>
340	Aug. 25, 1909	7.0	Dense.....	16.4	26.4	0.0	218
341	do.....	7.0	do.....	11.7	30.0	4.2	258
342	do.....	5.0	do.....	21.9	38.5	2.3	343
347	do.....	11.0	do.....	5.1	20.9	2.0	154
348	do.....	10.0	do.....	10.6	36.6	.6	247
349	do.....	5.0	do.....	20.3	57.7	.0	398
351	do.....	5.5	do.....	13.3	14.0	1.3	153
352	do.....	5.5	do.....	9.7	15.3	.3	130
438	Aug. 31, 1909	7.0	do.....	15.7	21.3	.7	195
439	do.....	8.0	do.....	11.1	43.0	12.3	408
440	do.....	6.0	do.....	15.4	23.6	.4	203
343	Aug. 25, 1909	14.0	Scattering.....	3.1	11.4	4.0	117
442	Aug. 31, 1909	12.0	do.....	5.8	19.3	.3	131
337	Aug. 25, 1909	11.0	Very scattering.....	.8	2.5	2.9	48
344	do.....	17.0	do.....	6.1	11.2	3.1	121
437	Aug. 31, 1909	14.0	Depleted.....	.0	1.8	.9	19
441	do.....	9.0	do.....	.0	1.4	1.4	22

POINT OF SHOALS ROCK.

This name is here given to a large area of varying productiveness lying between the preceding, Long Shoal and Point of Shoals Light-house, but it is possible that the name as used by the oystermen may not strictly accord with this usage. Scattered over the area are a number of small shoals ebbing nearly or quite bare, but the average depth is in general between 6 and 8 feet. Excepting at its northern edge, where an imaginary line separates it from Long Shoal Rock,

the bed is everywhere bounded by the deep water of the ship channel or a swash channel which separates it from Dry Shoals and Swash Rocks. Where it faces the ship channel there is for most of the distance a border of barren bottom lying between the bed proper and deep water.

The condition and extent of the bed at the time of the survey was as follows:

OYSTER GROWTH ON POINT OF SHOALS ROCK.

Character of growth.	Area.	Oysters per acre.	Total content of oysters.
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	254	200	50,800
Scattering.....	155	93	14,415
Very scattering.....	239	42	10,038
Depleted.....	142	15	2,130
Total.....	790	77,383

The dense areas are three, one near the eastern end of the bed, another adjoining the corresponding area of Long Shoal Rock, and the third an isolated spot on a shoal in the swash channel. The densest growths occur as a rule on the shoaler spots, especially at the eastern end of the bed, from the isolated area above alluded to to the ship channel. In this area the average growth is about 275 bushels of oysters to the acre—considering the depths, sufficient to yield about 38 bushels per day's tonging—while the average of the whole area of dense growth would not exceed 25 bushels per day.

There are four scattering areas, one of which, near the eastern apex of the bed, is insignificant. On these as a whole a tonger should average, at the beginning of the season, about 10 bushels per day.

The very scattering growth is distributed in three areas, of which one adjoins the dense growth on the isolated patch in the swash channel. They are barely prolific enough to raise them above the assumed limit of depletion.

The depleted area is in five patches or borders along the free boundary of the bed. They are entirely negligible in their productiveness.

On the dense and scattering areas the proportion of very small to small oysters is higher than on the beds previously described, and there are several places on each where the growth of oysters above 3 inches long is fair.

On the dense areas the deposit of shells is abundant, on the areas of scattering growth it is ample, while the areas with a very scattering growth and the depleted bottoms are decidedly deficient. In general the latter two areas are of no present and little prospective value.

DETAILS OF EXAMINATION OF POINT OF SHOALS ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.
				Spat.	Culls.	Counts.	
		<i>Feet.</i>					<i>Bushels.</i>
321	Aug. 25, 1909	5.0	Dense.....	8.7	10.5	0.3	101
327	do.....	5.5	do.....	11.9	10.3	.3	116
328	do.....	7.5	do.....	7.7	9.3	7.0	162
329	do.....	7.5	do.....	17.3	23.3	4.3	253
334	do.....	8.5	do.....	8.7	33.4	1.3	229
336	do.....	4.0	do.....	33.9	34.8	.3	353
356	do.....	10.0	do.....	4.3	16.0	5.7	165
311	Aug. 24, 1909	7.5	Scattering.....	8.0	6.3	.7	81
319	Aug. 25, 1909	7.0	do.....	.2	5.7	4.3	76
335	do.....	7.5	do.....	6.2	13.3	.4	104
354	do.....	8.0	do.....	.8	5.0	5.8	92
358	do.....	8.0	do.....	2.0	6.3	6.7	114
308	Aug. 24, 1909	7.5	Very scattering.....	.0	1.0	5.6	65
310	do.....	7.5	do.....	.3	.0	.3	5
320	Aug. 25, 1909	7.0	do.....	.7	2.3	1.7	34
357	do.....	8.0	do.....	1.3	2.7	2.7	50
527	Sept. 11, 1909	6.5	do.....	.5	4.0	2.9	54
309	Aug. 24, 1909	7.5	Depleted.....	.0	.0	.0	0
326	Aug. 25, 1909	9.5	do.....	.0	.0	1.1	12
333	do.....	10.5	do.....	1.7	2.1	1.7	38
355	do.....	9.0	do.....	.0	.0	.0	0
531	Sept. 11, 1909	7.0	do.....	.0	2.4	1.1	24

SWASH ROCK.

This bed lies inshore of the preceding, nearly surrounded by swash channels. It is connected by narrow isthmuses with Long Shoal and Dry Shoal Rocks and adjoins V Rock to the westward. It consists of a dense area surrounding two shoals ebbing bare, and two depleted areas which connect it with adjoining beds.

Its condition and extent in August, 1909, was as follows:

OYSTER GROWTH ON SWASH ROCK.

Character of growth.	Area.	Oysters per acre.	Estimated total content of oysters.
Dense.....	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Depleted.....	146	293	42,778
	115	15	1,725
Total.....	261		44,503

The dense area as a whole is prolific in its product, but the middle part of the bed, between the shoal spots, is less densely covered than the surrounding area. It is estimated that at the beginning of the season a tonger could take an average of about 39 bushels of oysters per day. The covering of clean shells is sufficient to guarantee a good strike under favorable conditions. The depleted bottom is practically bare of oysters and shells and is at present and potentially worthless under natural conditions.

The following examinations were made during the survey:

DETAILS OF EXAMINATION OF SWASH ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.
				Spat.	Culls.	Counts.	
324	Aug. 25, 1909	<i>Fect.</i> 5.0	Dense.....	29.2	41.4	0.0	<i>Bushels.</i> 360
325	do.....	7.0	do.....	12.3	17.3	7.0	226
228	Aug. 20, 1909	12.0	Depleted.....	.0	.0	.0	0
323	Aug. 25, 1909	9.0	do.....	.0	1.2	1.9	27
437	Aug. 31, 1909	14.0	do.....	.0	1.8	.9	19

MULBERRY SWASH ROCK.

This is a long narrow bed lying between Swash and V rocks on the outside and the so-called Marshy Island Rock on the shoreward side. At its southeastern end it is connected by narrow strips of indifferent productiveness with Wreck Shoal and Dry Shoal rocks, and its offshore boundary is the edge of the deep swash channel running toward Mulberry Point.

It consists essentially of bottom carrying a dense growth, interrupted at two places by areas of inferior productiveness. Its condition and extent in the latter part of August, 1909, are shown in the following table:

OYSTER GROWTH ON MULBERRY SWASH ROCK.

Character of growth.	Area.	Oysters per acre.	Estimated total content of oysters.
Dense.....	<i>Acres.</i> 422	<i>Bushels.</i> 302	<i>Bushels.</i> 127,444
Scattering.....	34	106	3,604
Very scattering.....	20	130	2,600
Depleted.....	29	43	1,247
Total.....	505	134,895

The dense area bears a growth varying from 161 to 570 bushels per acre, and the depths vary from 8 to upward of 20 feet. The heavier growth is as a rule in the shoaler water, though this rule is not without exceptions. It is estimated that a tonger could take an average of about 23 bushels per day at the beginning of the season.

The scattered area is limited in extent and bears a growth of between 88 and 117 bushels per acre, in a depth of between 13 and 15 feet, and it is estimated that it will yield about 8 bushels per day.

The area of very scattering growth connects this bed with Wreck Shoal Rock, and although, as shown by the foregoing table, the growth is heavier than on the preceding area, it lies in between 18 and 20 feet of water and will therefore be less productive to the tonger, its estimated initial yield being about 7 bushels per day. The depleted area is in several small patches.

There is an abundant supply of shells on the dense area, a quantity of doubtful sufficiency on the bottoms bearing scattering and very scattering growths, and a deficiency on the depleted bottom.

DETAILS OF EXAMINATION OF MULBERRY SWASH ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.
				Spat.	Culls.	Counts.	
		<i>Fect.</i>					<i>Bushels.</i>
213	Aug. 19, 1909	14.0	Dense.....	7.8	13.3	7.4	187
219	Aug. 20, 1909	10.0do.....	48.8	42.1	3.3	498
220do.....	12.0do.....	13.9	11.6	2.9	161
226do.....	17.5do.....	20.5	23.7	8.7	319
227do.....	12.5do.....	3.2	27.3	9.6	259
230do.....	12.0do.....	16.1	16.5	.6	173
238do.....	15.0do.....	26.5	31.9	2.3	322
331	Aug. 25, 1909	16.5do.....	16.8	30.0	6.8	312
338do.....	9.0do.....	23.8	80.0	4.2	570
346do.....	11.0do.....	28.9	34.0	.3	324
432	Aug. 31, 1909	16.5do.....	27.2	17.2	2.3	200
433do.....	13.0	Scattering.....	5.7	10.9	.3	88
435do.....	13.0do.....	5.8	10.7	2.7	113
436do.....	14.0do.....	3.6	17.3	1.0	117
345	Aug. 25, 1909	18.0	Very scattering.....	5.4	16.4	1.8	130
434	Aug. 31, 1909	17.0	Depleted.....	.5	4.1	1.8	43

MARSHY ISLAND ROCK.

This lies between Mulberry Swash Rock and the inshore boundary of the public ground, principally in the "addition" which was made a part of the ground subsequent to the Baylor survey. The name here employed is coined for the purpose of this report, as the name by which this area of oyster bottom is known to the oystermen, if it has a distinctive name, was not ascertained by the survey. The outer or offshore boundary of the bed is defined more or less sharply by a channel, carrying a maximum of from 21 to 27 feet of water, between this and Mulberry Swash Rock.

The condition and extent of this bed about the middle of August, 1909, was as follows:

OYSTER GROWTH ON MARSHY ISLAND ROCK.

Character of growth.	Area.	Oysters per acre.	Estimated total content of oysters.
Dense.....	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Scattering.....	197	231	45,507
Very scattering.....	322	129	41,538
Depleted.....	235	85	19,975
	387	18	6,966
Total.....	1,141		113,986

The dense areas lie in three isolated patches which exhibit no material shoaling over the surrounding bottom, except where they touch the channel which bounds the bed offshore. The depth of water on

the several areas ranges between 6 and 16 feet, and the density of growth between 142 and 410 bushels per acre. It is estimated that the bottoms are capable of producing an initial yield of about 18 bushels per day.

The scattering area occupies a general central position in the bed surrounding one of the dense spots. The depth varies from 10 feet inshore to about 22 feet at the edge of the channel, and the quantity of oysters varies between 92 and 186 bushels per acre. The estimated daily yield to the tonger is about 10 bushels.

The bottom covered by a very scattering growth forms a zone encircling the inner edge of the preceding. It lies in a depth varying from 7 to 16 feet, and, although the examinations were not as numerous as they should have been, they indicate that the growth is sufficient to yield an average of between 6 and 7 bushels per day.

The depleted bottom lies in a belt on the inside edge of the bed. It is practically bare of oysters and shells.

Shells are found in fair quantities on the dense bottom and on the outer parts of the scattering growth, but are deficient on the inshore parts of the latter, on most of the area of very scattering growth, and on the depleted area.

The following observations were made:

DETAILS OF EXAMINATION OF MARSHY ISLAND ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.
				Spat.	Culls.	Counts.	
		<i>Feet.</i>					<i>Bushels.</i>
217	Aug. 19, 1909	13.0	Dense	5.4	12.7	4.6	142
225	Aug. 20, 1909	12.5	do	6.2	23.1	1.9	170
232	do	14.0	do	10.4	54.3	8.3	410
239	do	11.5	do	7.7	11.5	9.6	201
221	do	13.0	Scattering	6.7	9.6	.8	92
222	do	14.0	do	8.1	9.2	1.2	102
224	do	9.5	do	2.1	8.8	7.5	136
231	do	17.0	do	10.9	23.6	.9	186
233	do	11.0	do	7.3	12.1	2.8	129
223	do	9.0	Very scattering	1.2	5.8	4.2	81
234	do	16.0	do	4.5	15.9	.5	109
218	Aug. 19, 1909	12.0	Depleted0	.8	1.3	18

LONG SHOAL ROCK.

This triangular bed flanks a shoal, ebbing bare in many places, which extends eastwardly from Point of Shoals light-house for a distance of upward of $1\frac{1}{2}$ miles. As understood in this report, its boundary is an imaginary line running from Point of Shoals light toward Jail Island at an average distance of about 300 to 400 yards from the crest of the shoal, as far as the swash channel opening toward the northwest, along the edge of this channel to its mouth, and thence to the starting point. The main body of the rock, therefore, lies north of the crest of the bar. Its condition and extent about the beginning of September, 1909, were as follows:

OYSTER GROWTH ON LONG SHOAL ROCK.

Character of growth.	Area.		Estimated total content of oysters.
	<i>Acres.</i>	<i>Bushels.</i>	
Dense.....	331	241	79,771
Scattering.....	10	64	640
Very scattering.....	84	60	5,040
Depleted.....	79	16	1,264
Total.....	504		86,715

A dense growth of small oysters, with a good proportion of larger ones in a few spots, covers the major part of the bed. At various places the total growth varies between 148 and 364 bushels per acre, and it is estimated that at the beginning of the season a tonger could take about 28 bushels per day.

The scattering growth is comprised in one small spot about 100 to 200 yards from the light-house, where about 9 bushels per day may be taken.

The very scattering growth lies in two small patches along the western border of the bed and a larger area south of the ridge is continuous with similar bottom on Point of Shoals Rock. It is capable of yielding between 6 and 7 bushels per day. The depleted area lies south of the outer half of the ridge, with a small patch on the swash channel. It is practically bare of oysters and shells.

The areas bearing oysters in dense and scattering growth are covered with a supply of shells amply sufficient to serve the purposes of culch. The small patches of very scattering growth are also fairly covered, but the large area south of the ridge and the depleted area adjoining are deficient.

The following observations were made:

DETAILS OF EXAMINATION OF LONG SHOAL ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.
				Spat.	Culls.	Counts.	
		<i>Feet.</i>					<i>Bushels.</i>
295	Aug. 24, 1909	8.0	Dense.....	25.8	29.7	2.3	308
296	do.	10.0	do.	20.4	25.4	.4	237
298	do.	12.5	do.	11.9	13.4	9.6	232
299	do.	6.5	do.	22.8	17.1	2.5	230
300	do.	6.5	do.	19.2	10.0	12.5	283
301	do.	7.5	do.	20.3	7.3	.7	148
313	do.	8.0	do.	19.7	25.8	.0	232
318	do.	6.0	do.	40.7	30.7	.0	364
322	Aug. 25, 1909	5.0	do.	.0	4.6	13.1	164
529	Sept. 11, 1909	5.0	do.	19.7	21.4	.0	210
541	do.	6.0	Scattering	1.1	6.6	2.3	64
297	Aug. 24, 1909	11.0	Very scattering	6.1	4.5	1.9	74
312	do.	9.0	do.	2.1	3.3	4.2	72
538	Sept. 11, 1909	4.0	do.	3.9	3.2	.0	36
307	Aug. 24, 1909	8.5	Depleted.....	1.4	1.1	2.5	39
528	Sept. 11, 1909	6.5	do.	.0	1.1	.5	11
554	do.	7.0	do.	1.3	3.4	.3	27
535	do.	7.0	do.	.0	.5	.3	6
539	do.	6.0	do.	.2	.5	.2	6
550	do.	6.0	do.	.5	.5	.2	7

V ROCK.

This bed takes its name from the shape of a bare shoal near its southwestern edge. It is inshore of the preceding rock and adjoins Swash Rock to the southeast.

The area and character of growth on the bed are epitomized in the following table:

OYSTER GROWTH ON V ROCK.

Character of growth.	Area.		Estimated total content of oysters.
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	240	227	54,480
Very scattering.....	73	84	6,132
Depleted.....	73	15	1,095
Total.....	386	61,707

The dense area occupies the middle belt of the bed and carries a growth of between 144 and 344 bushels per acre, the average estimated yield per day to the tonger being about 21 bushels.

The very scattering growth lies along the northwestern edge of the bed and on a comparatively shallow ridge along the swash channel near its mouth. It bears oysters in a quantity to yield about 7 bushels per acre.

The depleted area adjoins similar bottom on Swash Rock and is practically bare of both oysters and shells. The supply of shells on the rest of the bed is ample to secure their reseeding under proper conditions.

The data for the bed are as follows:

DETAILS OF EXAMINATION OF V ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.
				Spat.	Culls.	Counts.	
		<i>Feet.</i>					<i>Bushels.</i>
229	Aug. 20, 1909	9.0	Dense.....	31.8	16.5	1.7	265
235do.....	12.5do.....	10.8	45.7	.4	293
236do.....	10.5do.....	20.6	19.4	.0	204
268	Aug. 23, 1909	12.5do.....	4.4	15.2	9.6	203
303	Aug. 24, 1909	9.5do.....	21.7	10.4	.8	173
304do.....	7.5do.....	18.3	10.0	.0	144
305do.....	6.5do.....	47.7	19.7	.0	344
306do.....	15.5do.....	13.9	14.6	4.8	197
317do.....	10.0do.....	22.5	18.7	.8	219
302do.....	9.5	Very scattering.....	2.5	7.9	2.9	84
315do.....	10.0	Depleted.....	1.3	.4	2.1	31
316do.....	10.0do.....	.0	.0	.0	0

MOORES ROCK.

The bed known to the oystermen by this name lies on a shoal surrounded by deep water about halfway between Point of Shoals Light-House and Mulberry Point. It consists principally of bottom bearing a dense growth, with a scattering fringe along the southern half of its western edge. Its general extent and condition are as follows:

OYSTER GROWTH ON MOORES ROCK.

Character of growth.	Area.	Oysters per acre.	Estimated total content of oysters.
Dense.....	<i>Acres.</i> 37	<i>Bushels.</i> 265	<i>Bushels.</i> 9,805
Scattering.....	6	168	1,008
Total.....	43	10,813

On the dense area the oysters, as developed by the survey, range between 134 and 351 bushels per acre, and it is estimated that the bottom as a whole will produce about 28 bushels of oysters per day of tonging.

The area of scattering growth lies in the deeper water close to the adjoining barren bottom, and its estimated yield to the tonger is about 8 bushels per day. The deposit of shells is good over the entire area of the bed.

DETAILS OF EXAMINATION OF MOORES ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.
				Spat.	Culls.	Counts.	
		<i>Feet.</i>					<i>Bushels.</i>
241	Aug. 20, 1909	9.5	Dense.....	33.4	35.4	0.0	351
265	Aug. 23, 1909	10.5do.....	19.6	19.2	2.5	225
266do.....	6.5do.....	28.3	33.4	3.3	350
267do.....	5.5do.....	13.3	13.0	.0	134
242	Aug. 20, 1909	20.5	Scattering.....	10.4	12.5	.0	168

HORSEHEAD ROCK.

This bed covers several shoals along the edge of deep water south of Mulberry Point, and for the purpose of this report is considered to include a small patch close to the Baylor line to the eastward. The apex of the bed is detached, but the remainder is continuous, though of varying productiveness. East of this rock and north of Marshy Island Rock the survey found small patches of oysters close to the Baylor line, adjoining or included in various planted beds. This region is shown on the charts, included within red lines but without shading.

The general distribution of oysters on Horsehead Rock is as follows:

OYSTER GROWTH ON HORSEHEAD ROCK.

Character of growth.	Area.		Estimated total content of oysters.
	<i>Acres.</i>	<i>Bushels.</i>	
Dense.....	33	223	7,359
Scattering.....	192	104	19,968
Very scattering.....	139	112	15,568
Depleted.....	16	0	0
Total.....	380		42,895

The areas of dense growth are on the terminal isolated shoal and in two small patches on the main part of the bed. The growth varies between 178 and 283 bushels per acre, and it is estimated that the area as a whole will yield an average of 20 bushels per day to the tonger at the beginning of the season.

On the areas of scattering oysters the density of growth is between 47 and 170 bushels per acre, and it is estimated that they are capable of yielding, at the beginning of the season, an average of about 7 bushels per day per tonger.

On the bottom which is rated as carrying a very scattering growth the average per acre is slightly higher than on the preceding, but as the water is deeper it is less productive in its return per day of labor expended on it.

The depleted bottom lies inshore, close to Mulberry Point, and is practically, in many cases absolutely, bare of oysters and almost as deficient in shells. On all other areas the deposit of shells is good or fair.

DETAILS OF EXAMINATION OF HORSEHEAD ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.
				Spat.	Culls.	Counts.	
		<i>Fect.</i>					<i>Bushels.</i>
256	Aug. 23, 1909	14.0	Dense.....	19.3	35.4	0.4	283
257	do.....	7.0	do.....	16.5	22.7	.3	203
258	do.....	16.0	do.....	6.5	29.6	4.1	227
263	do.....	7.0	do.....	9.1	24.5	.6	178
255	do.....	18.5	Scattering.....	4.1	6.4	10.9	170
259	do.....	9.0	do.....	6.5	16.9	1.2	122
261	do.....	12.0	do.....	5.4	15.1	.5	110
262	do.....	7.0	do.....	8.1	5.9	.3	74
264	do.....	12.0	do.....	7.1	12.5	.4	104
273	Aug. 24, 1909	4.5	do.....	.4	5.6	1.1	47
260	Aug. 23, 1909	16.0	Very scattering.....	4.5	16.3	.5	112
274	Aug. 24, 1909	6.0	Depleted.....	.0	.0	.3	4
280	do.....	5.5	do.....	.0	.2	.0	1
291	do.....	5.0	do.....	.0	.0	.0	0
292	do.....	4.5	do.....	.0	.0	.0	0
293	do.....	5.0	do.....	.0	.0	.0	0
294	do.....	7.5	do.....	.0	.0	.0	0

DEEPWATER SHOALS ROCK.

This is considered as including all oyster bottoms within the Baylor lines above Mulberry Point. Its condition and extent are as follows:

OYSTER GROWTH ON DEEPWATER SHOAL ROCK.

Character of growth.	Area.		Estimated total content of oysters.
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	17	129	2,193
Very scattering.....	21	57	1,097
Depleted.....	241	12	2,892
Total.....	279	6,182

The comparatively small productive area on this bed all lies within a radius of about 1,000 yards of Deepwater Shoals Light-House, most of it being in the immediate vicinity of the light. The dense area is in two small patches on which there is a sufficient growth to yield an average maximum of about 15 bushels per day of actual tonging.

The very scattering areas are three in number, all more or less intimately associated with the preceding. They should yield about 6 bushels per day at the beginning of the season.

The depleted area is practically devoid of oysters. On the areas of dense and very scattering growth there is a good covering of shells, and they are also found in ample numbers on the depleted area within a radius of 1,000 or 1,200 yards of the light, but elsewhere the bed is practically denuded.

The following examinations were made:

DETAILS OF EXAMINATION OF DEEPWATER SHOALS ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.
				Spat.	Culls.	Counts.	
		<i>Feet.</i>					<i>Bushels.</i>
269	Aug. 23, 1909	4.0	Dense.....	4.6	12.6	0.3	91
270	do.....	9.0	do.....	2.1	12.9	5.8	138
285	Aug. 24, 1909	8.0	do.....	1.2	17.0	.9	158
272	Aug. 23, 1909	5.0	Very scattering.....	1.2	7.3	.0	44
284	Aug. 24, 1909	10.0	do.....	.0	.8	4.2	49
286	do.....	8.0	do.....	.0	7.7	3.3	74
288	do.....	7.0	do.....	.9	4.9	3.0	62
271	Aug. 23, 1909	6.0	Depleted.....	.3	2.9	.6	23
282	Aug. 24, 1909	5.0	do.....	.0	.2	.0	1
283	do.....	6.0	do.....	.3	1.5	.0	9
287	do.....	7.0	do.....	.7	2.2	.0	15

ROCK WHARF SHOALS ROCK.

This bed lies near the western end of Public Ground No. 1, Isle of Wight County, across the river from the group of seed beds previously described. It forms two patches surrounding shoals and consists principally of productive bottom, as shown in the following table:

OYSTER GROWTH ON ROCK WHARF SHOALS ROCK.

Character of growth.	Area.	Oysters per acre.	Estimated total content of oysters.
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	18	140	2,520
Depleted.....	8	11	88
Total.....	26		2,608

The dense area should yield an average of about 22 bushels of oysters per day, and is fairly covered with clean shells. The depleted bottom is practically bare of both oysters and shells.

DETAILS OF EXAMINATION OF ROCK WHARF SHOALS ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.
				Spat.	Culls.	Counts.	
		<i>Feet.</i>					<i>Bushels.</i>
421	Aug. 28, 1909	4	Dense.....	7.3	13.6	2.1	129
422do.....	6do.....	11.3	15.7	1.4	152
420do.....	3	Depleted.....	.0	.2	.9	11

BEDS BETWEEN ROCK WHARF SHOALS AND SPINDLE ROCK.

These cover the largest area of productive bottom in the ground, distributed in three patches. Their aggregate area and extent are as follows:

OYSTER GROWTH ON BEDS BETWEEN ROCK WHARF SHOALS AND SPINDLE ROCK.

Character of growth.	Area.	Oysters per acre.	Total content of oysters.
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	45	140	6,300
Very scattering.....	5	46	230
Depleted.....	37	18	666
Total.....	87		7,196

The dense areas are close to the shoal spots and in various places bear from 101 to 178 bushels per acre, the average density being sufficient to yield about 22 bushels per day to the tonger.

The bottom of very scattering growth covers but a small spot at the outer end of the middle shoal, and the density of growth is sufficient to yield barely 5 bushels of seed oysters per day.

The depleted bottom is practically denuded. It bears very few shells, and the very scattering bottom is little better in this respect, but the shell deposit on the dense areas is good.

DETAILS OF EXAMINATION OF BEDS BETWEEN ROCK WHARF SHOALS AND SPINDLE ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.
				Spat.	Culls.	Counts.	
		<i>Feet.</i>					<i>Bushels.</i>
409	Aug. 28, 1909	4.0	Dense.....	8.6	18.2	1.0	147
417do.....	6.0do.....	17.0	10.0	.8	146
418do.....	7.0do.....	9.2	9.0	2.6	121
419do.....	5.0do.....	9.4	20.0	2.6	178
425do.....	3.5do.....	6.2	8.4	2.5	101
424do.....	7.5	Very scattering.....	.5	3.1	2.6	46
410do.....	6.5	Depleted.....	.3	1.3	1.3	22
416do.....	7.0do.....	.0	.3	1.3	15

SPINDLE ROCK.

This bed follows the line of a shoal at right angles to the shores. It consists principally of a dense growth, with insignificant areas of very scattering oysters and depleted bottom at its inner end. Its area and condition at the time of the survey were as follows:

OYSTER GROWTH ON SPINDLE ROCK.

Character of growth.	Area.	Oysters per acre.	Estimated total content of oysters.
	<i>Acres.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Dense.....	14	140	1,960
Very scattering.....	3	27	81
Depleted.....	2	12	24
Total.....	19	2,065

The dense area bears a growth of between 119 and 179 bushels per acre, and is capable of producing about 21 bushels of oysters per day's tonging; the area of very scattering growth will yield barely 5 bushels and the depleted bottom about 2 bushels. The area of dense growth bears a good supply of shells, that of very scattering growth hardly enough to insure reseeding except under the best conditions, while the depleted bottom is deficient.

The following examinations were made:

DETAILS OF EXAMINATION OF SPINDLE ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.
				Spat.	Culls.	Counts.	
415	Aug. 28, 1909	<i>Fect.</i> 6.0	Dense.....	7.7	19.7	3.7	<i>Bushels.</i> 179
426	do.	4.0	do.	8.8	14.0	.4	120
427	do.	6.0	do.	7.4	13.6	1.1	119
411	do.	4.0	Very scattering.....	1.3	2.6	.7	27
412	do.	4.5	Depleted.....	.0	.9	.7	12

DAYS POINT SHOAL BED.

This follows a shoal but part of which is included in the public ground. The part included embraces a dense growth capable of yielding to the tonger about 27 bushels of oysters per day. The following is the result of the examination made:

DETAILS OF EXAMINATION OF DAYS POINT SHOAL ROCK.

Station number.	Date of examination.	Mean depth of water.	Character of growth.	Oysters caught per square yard.			Estimated quantity oysters per acre.
				Spat.	Culls.	Counts.	
413	Aug. 28, 1909	<i>Fect.</i> 4.5	Dense.....	7.2	22.3	1.5	<i>Bushels.</i> 166

PUBLIC GROUNDS.

The public oyster grounds of Virginia are those areas of the bottoms of tide water which are included within the lines of the Baylor survey and additions thereto upon which the public is permitted to take oysters at certain seasons of the year on compliance with certain conditions, and which are withheld from lease for purposes of oyster culture under private and exclusive control.

The public grounds were designed to include all of the natural rocks, though, as has been explained previously, no actual examination was made for the purpose of really determining the facts. The boundaries are necessarily straight lines and do not purport to conform to the outlines of the actual rocks, and largely for this reason they can not fail to include within their confines more or less barren bottom. The relation which the barren bottoms bear to that which actually produces oysters has been in more or less acrimonious dispute between the tongers and dredgers on the one hand and the planters and their partisans on the other, and it was largely to secure authentic and definite information on this point that the present survey was undertaken.

The public grounds are officially designated by numbers and the name of the county within which they are supposed to lie, and on the accompanying charts their boundaries, accurately platted from the charts of the Baylor survey, are indicated by broken black lines. The boundaries of the natural rocks, as determined by the present survey, are shown in solid red lines, within which the varying density of oyster growth is shown by the relative density of the shading. An inspection of the charts will show that the natural rocks are more or less scattered, between and about them lying barren bottoms, shown as unshaded areas, within the boundaries of the Baylor survey.

At various places it was found that certain private grounds, as indicated by the boundary stakes, encroached more or less on the public grounds, though from the flimsy character of the marks it was difficult in many cases to determine the real facts. This apparent encroachment of private interests on the public domain was observed at various places in Nansemond River, between Fishing Point and Ballards Marsh, about Creek Channel Shoal and Aaron Shoal rocks, in the vicinity of Browns Shoal rocks, at the inshore edges of Kettle Hole and Blunt Point rocks, and at various places between Jail Island and Mulberry Point.

Whatever may have been the conditions under which this encroachment was originally permitted, it was undoubtedly aided by the latter-day uncertainty as to the Baylor boundaries. Apparently but little effort has been made to maintain or replace the shore marks to which the corners of the Baylor survey were referred, and a number of them appear to be now unavailable for reference. The irregularity of the boundaries has also made the maintenance of the lines more difficult, and the same conditions have made it almost impossible for the oyster police to prevent the planters from degrading the public beds beyond their staked boundaries.

These reasons have made it important to both "natural growers" and planters that an examination should be made into the actual location of the productive areas or those which, though at present more or less unproductive, may be reasonably expected to recuperate under proper natural conditions.

To assist to an understanding of the conditions on the public beds as a whole the following discussion is offered. The several public beds in the region surveyed are considered with regard to the relative areas of dense, scattered, very scattered, and depleted growths, and barren bottom. The first four are measured from the results of the present survey, while the barren bottom is regarded as the difference between the sum of these areas and the areas of the public beds according to Baylor's computations, the data being exhibited in tabular form for each of the several public grounds. For each public ground or for each fraction or combination considered as an entity in the following pages, there are furnished tables and

summaries of the estimated total contents of market oysters, as distributed by rocks and varying densities of growth. These estimates are interesting, but are misleading if regarded as a measure of productiveness, for a very sparse growth over a large area, as compared with a dense growth over a small one, will give a great aggregate which really represents nothing commercially, as the oysters may be so thinly scattered as to be totally unavailable industrially.

The important point is not how many oysters there may be on a given bed at a given time, but the quantity of oysters available under existing local economic conditions, the maximum number of bushels that can be removed with profit to the tonger.

It is unnecessary to explain to those familiar with the oyster industry that it is practically impossible to accomplish a complete denudation of the beds in any one season, but there are cases known to the writer, though he has no personal knowledge of the kind in the region under discussion, in which small rocks have been, in effect, taken up bodily, oysters, seed, and shells, and transferred to planted beds.

Under ordinary circumstances, in localities where the cull laws can be and are reasonably enforced, not only the seed or young oysters but a considerable proportion of the market oysters are left on the beds at the end of the season. Eventually, however, the oysters become so scattered that the daily yield to the tonger becomes less than a minimum daily wage, and while the aggregate quantity of marketable oysters left on the beds appears large when expressed in a total of bushels, as in the tables of total contents, it will no longer pay to take them. The minimum average density of growth to which a bed may be reduced before becoming commercially unproductive depends primarily upon the price of oysters. The smaller the market value of a bushel of oysters the greater is the quantity that must be taken per day to furnish a living wage. Another factor that is essentially involved is the amount of culling required, less labor being necessary in handling the oysters when they are single or in small clusters than when they are badly clustered and overgrown with young, from which they must be separated before being placed on the market.

The depth of water is also a very important factor in determining the actual density of growth necessary to render a bed commercially productive. As has been explained in describing the methods pursued in the preparation of this report, the deeper the water the greater must be the quantity of oysters per square yard or acre necessary to afford the tonger a given catch per day. Not only do his tongs of any given length of shaft and head cover a smaller area on the bottom, but the time and labor of making the "grab"—that is, putting the tongs on the bottom, scraping up the oysters, and pulling them up—are materially increased. In other words, in deep water

not only is the area covered by a "grab" smaller, but, other things being equal, fewer "grabs" can be made in an hour than in a smaller depth.

In the tables shown in this report and on the chart these factors have all been considered in estimating the relative density of the beds. In the estimation of the available contents of the rocks as exhibited in the following tables the same factors have been considered. It is assumed that, at the price which has recently been received for market oysters in the region under consideration—namely, 45 cents per bushel—it would be wholly unprofitable to tong on bottoms which would yield less than 3 bushels of culled oysters per day, exclusive of the time spent in culling, which would ordinarily involve part of the time of a second man or boy. In the same way at the price of seed oysters, namely 30 cents per bushel, it is assumed to be equally unprofitable to tong on bottoms yielding less than 4 bushels, exclusive of shells.

It can not be argued that this limit is too high, but undoubtedly it will be claimed by some that it is entirely too low. The objection would be well founded if it were to apply wholly to areas on which the initial density of growth was such as to afford the minimum yield adopted, but it will not lie against the application of the standard to areas of greater initial productiveness. A dense bed in course of partial denudation by tonging is not uniformly depleted over its whole area. The tongers spread themselves more or less promiscuously over the rocks and take up practically all of the oysters in patches, while other areas are, for the time being, inadvertently left untouched. Later many of these untouched spots are tonged with profit, until the worked areas become so great in proportion to those which have been overlooked that the time spent in searching for the latter makes further work unremunerative. At this stage of temporary abandonment the rock consists of a few small patches of productive bottom, areas which are practically bare of market oysters, and others which have been worked over but still retain some oysters scattered over them by the operations of tonging. It is of course impossible, from the complexity and irregularity of the conditions obtaining on an oyster bed, to fix a limit of more than reasonable accuracy. In preparing the following tables the present available productiveness of each area has been considered with regard to the terms of its initial yield to the tonger and its total estimated contents above that which would give a return of 3 bushels per day's work on the market oyster beds and 4 bushels on the seed beds. The depleted areas and most of the areas covered by what is called very scattered growth are therefore negligible as present factors. A very few areas in the depleted bottoms and a somewhat greater proportion of the bottoms bearing a very scattering growth are of potential value as bearing small oysters and shells which reasonably assure future regeneration.

These are pointed out in the following discussion of the several public grounds:

PUBLIC GROUNDS NO. 2 NANSEMOND COUNTY AND NO. 6 ISLE OF WIGHT COUNTY.

These two grounds overlap, as platted on the state charts, and as they can not be accurately differentiated in the conflicting area they may be most conveniently considered together. The former bed begins at the upper limit of oyster growth in the Nansemond River off Cedar Point, and becoming continuous with No. 6 near Newport News Rock, the latter extends along the right side of the James River to beyond Ballards Marsh Rock. Ground No. 2 is said to contain 3,319.6 acres, and Ground No. 6, 4,148.2 acres, a total of 7,467.8; but there is an overlap or duplication of about 305 acres, and deducting this, the actual total area of the two beds may be assumed to be about 7,162.8 acres. The following is a résumé of the extent of the oyster bottoms of the several rocks and the barren bottom embraced within the limits of these grounds:

AREAS OF OYSTER GROWTH IN PUBLIC GROUNDS NO. 2 NANSEMOND COUNTY AND NO. 6 ISLE OF WIGHT COUNTY.

Name of oyster rock.	Oyster growth.				
	Dense.	Scatter- ing.	Very scatter- ing.	Depleted.	Total.
	Acres.	Acres.	Acres.	Acres.	Acres.
Larkins.....	0	0	0	39	39
Nansemond Ridge.....	85	446	294	782	1,607
Drum Shoal.....	0	19	14	95	128
Newport News.....	4	27	12	129	172
Cruiser Shoal.....	27	19	26	32	104
Between Nansemond Ridge and Fishing Point ^a	50	0	5	7	62
High shoal.....	24	13	24	95	156
Trout Shoal.....	0	25	14	90	129
Dog Shoal.....	16	11	35	120	182
Fishing Point.....	45	77	47	90	259
Between Fishing Point and Ballards Marsh.....	5	0	8	18	31
Ballards Marsh.....	0	4	33	142	179
Total oyster area.....	256	641	512	1,639	3,048
Total barren bottom.....					4,114.8
Total Baylor survey.....					7,162.8

^a 8 acres undetermined.

It will be observed from this table that the barren bottom, as developed by this survey, exceeds the area of the oyster rocks and constitutes about 57 per cent of the area of the two public grounds under discussion. The depleted bottom, which, excepting the places noted in the detailed descriptions of the several beds, is at present unproductive and of a character that gives little or no promise of future regeneration, forms about 23 per cent of the total area included within the Baylor lines. Assuming that the areas of very scattering

growth, at present practically worthless so far as actual productiveness is concerned, are capable of coming into production at some time in the future, by virtue of the young growth and clean shells that they bear, it will be seen that the actual productive oyster rocks form only about 20 per cent of the area of these two public grounds.

The estimated total marketable contents of the grounds, based upon the distribution of oysters as indicated by the chain, and the actual productiveness of the various areas as determined by actual count and measurement, is exhibited in the following table:

CONTENT OF MARKET OYSTERS, PUBLIC GROUNDS No. 2 NANSEMOND-COUNTY AND No. 6 ISLE OF WIGHT COUNTY.

Name of rock.	Dense.	Scatter- ing.	Very scatter- ing.	Depleted.	Total.
	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Larkins.....				195	195
Nansemond Ridge.....	7,905	26,760	10,878	8,602	54,145
Drum.....		950	546	2,755	4,251
Newport News.....	432	1,701	420	3,483	6,036
Cruiser Shoal.....	4,212	1,007	728	288	6,235
Flat Rock, etc.....	5,400		210	182	5,792
High Shoal.....	2,160	624	600	807	4,191
Trout Shoal.....		1,100	420	720	2,240
Dog Shoal.....	1,664	507	945	1,416	4,532
Fishing Point.....	5,355	6,314	2,068	1,710	15,447
Between Fishing Point and Ballards Marsh.....	1,340		248		1,588
Ballards Marsh.....		124	792	894	1,810
Total.....	28,468	39,087	17,855	21,052	106,462

This indicates that if it were possible to "clean up" completely the entire area covered by the oyster rocks, the product would be about 106,000 bushels of marketable oysters. When an analysis is made, it is speedily apparent that the commercially available supply on these beds is only about 40 per cent of the foregoing, as stated in the following table:

AVAILABLE CONTENT OF MARKET OYSTERS, PUBLIC GROUNDS No. 2 NANSEMOND COUNTY AND No. 6 ISLE OF WIGHT COUNTY.

Name of rock.	Dense.	Scatter- ing.	Very scatter- ing.	Total.
	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Larkins.....				
Nansemond Ridge.....	5,500	13,500	1,000	20,000
Drum.....		400	100	500
Newport News.....	350	850		1,200
Cruiser Shoal.....	3,400	700		4,100
Flat Rock, etc.....	3,600			3,600
High Shoal.....	1,500	350	100	1,950
Trout Shoal.....		600		600
Dog Shoal.....	1,200	250		1,450
Fishing Point.....	4,400	3,600	300	8,300
Between Fishing Point and Ballards Marsh.....	1,200			1,200
Ballards Marsh.....		75	100	175
Total.....	21,150	20,325	1,600	43,075

It will be observed that on the basis assumed in this report the depleted areas are wholly unproductive commercially, and the bottoms covered with very scattered growth are practically so. On the latter the growth in many cases is barely sufficient to yield 3 bushels per day, and in no case does it much exceed that limit. The large aggregate of market oysters on the areas of very scattered and depleted bottoms are so thinly distributed as to be unavailable commercially, and are therefore valueless except as brood stock to assist in furnishing spat for replenishing the beds. On the dense areas about three-fourths of the total contents and on the scattering growths about one-half may be taken with profit.

The total estimated available product of 43,075 bushels appears very small as compared with the area included within the Baylor lines, averaging but about 6 bushels per acre. It is about half of the average yield of marketable oysters on the public grounds of the State as a whole in 1901 and 1904, according to the statistics of the Bureau of Fisheries, and about equal to the average yield in 1908, as stated by the Bureau of the Census.

The deficiency in productiveness of this section was to be expected in view of public report. The beds, especially in Nansemond River, are generally recognized as being seriously depleted, the allegation of the tongers being that several years ago large quantities of uncultured stock were taken from the beds for deposit on private planting ground, and the tonger employed by the survey is authority for the statement that the growth on the Nansemond River beds in the season preceding the investigation was hardly sufficient to warrant tonging.

Combining the exhibits of the tables of areas and of commercially available oysters, we find that it apparently would be profitable to take from the dense growths about 83 bushels per acre and from the scattering growths an average of about 32 bushels. On the bottoms with a very scattering growth the average content per acre at the beginning of the present oyster season was so small that, even under the very low standard of profit adopted in this report, the beds would be reduced to unproductiveness after an average of only about 3 bushels of oysters per acre had been removed. Of course a very large part of this bottom must be regarded as practically unproductive in the beginning, and it is only here and there that even the least ambitious tonger would venture to work.

Another aspect of the present state of these grounds is the production of young oysters and the presence of shells in such quantities and cleanliness as to afford prospect of a strike under proper conditions. The following table gives the estimated total content of the several rocks and of the grounds as a whole in oysters less than 3 inches long:

TOTAL CONTENT OF YOUNG OR SEED OYSTERS, PUBLIC GROUNDS No. 2 NANSEMOND COUNTY AND No. 6 ISLE OF WIGHT COUNTY.

Name of rock.	Dense.	Scatter- ing.	Very scatter- ing.	Depleted.	Total.
	<i>Bush.</i>	<i>Bush.</i>	<i>Bush.</i>	<i>Bush.</i>	<i>Bush.</i>
Larkins.....				390	390
Nansemond Ridge.....	12,580	59,318	9,996	27,370	109,264
Drum.....		1,748	854	5,890	8,492
Newport News.....	372	2,025	996	4,386	7,779
Cruiser Shoal.....	3,780	893	1,326	2,112	8,111
Flat Rock, etc.....	150		275	112	537
High Shoal.....	3,216	1,651	1,392	760	7,019
Trout Shoal.....		4,112	1,652	1,890	7,654
Dog Shoal.....	2,480	1,989	770	4,838	10,077
Fishing Point.....	8,325	13,706	3,290	2,700	28,021
Between Fishing Point and Ballards Marsh.....	915	0	0	0	915
Ballards Marsh.....		608	6,303	6,590	13,501
Total.....	31,818	86,050	26,854	57,038	201,700

In individuals the small oysters are five or six times as numerous as the market oysters and in measured quantity they are about twice as abundant. On the dense areas they bulk about the same as the market oysters, but as individuals they are two or three times as many. On the scattered area they much exceed the market oysters in numbers and are more than double them in measured quantity. As both of these types of bottom are almost invariably supplied with cultch in the form of clean shells, it can be safely assumed that their future is assured under ordinarily fair conditions and provided the beds are not stripped under infractions of the culling law.

On the area of very scattering growth the quantity of young in nearly every case materially exceeds that of market oysters. Almost the sole exception is Nansemond Ridge Rock, where the young and market oysters are about equal in quantity, the former being decidedly deficient in all places excepting close to the denser areas below a line between Pig and Barrel points.

Excepting Nansemond Ridge Rock the very scattering areas bear an average of about two and one-half times as many bushels of young as of old oysters per acre, and there is nearly everywhere a sufficient abundance of shells to justify the prediction of future regeneration if man will permit. On Nansemond Ridge the future of the very scattering areas, except in a few places, appears unpromising.

The depleted area is, on the whole, deficient in shells and young oysters, and if we except one or two spots near Nansemond Light, the outer end of Ballards Marsh Rock, and several other places quite close to the productive areas, there is but little probability that any of the area will become naturally productive.

PUBLIC GROUND NO. 1 WARWICK COUNTY, BELOW DEEP CREEK.

This public ground, while continuous in its lines from near Newport News to above Deepwater Shoals Light, is divided, for purposes of administration, by a line running from Deep Creek to Days Point. Below this line the cull law is in force and tonging is practically confined to taking oysters for the market, while above the line it is legally permissible to take oysters of all sizes for planting purposes. The total area of the portion of the bed here discussed is about 5,515 acres. It embraces six well-defined rocks or groups of rocks, the general condition and area of which are shown in the following table, which also includes a very small contiguous and overlapping area at the inshore edge of Kettle Hole Rock, known as Public Ground No. 2 Warwick County:

AREAS OF OYSTER GROWTH, PUBLIC GROUND NO. 1 WARWICK COUNTY, BELOW DEEP CREEK.

Name of oyster rock.	Dense.	Scatter- ing.	Very scat- tering.	Depleted.	Total.
	<i>Acres.</i>	<i>Acres.</i>	<i>Acres.</i>	<i>Acres.</i>	<i>Acres.</i>
Browns Shoal.....	68	44	27	226	365
Gun Shoal.....	6	0	16	4	26
Kettle Hole.....	258	66	111	11	446
Thomas Point.....	76	118	100	127	421
Blunt Point.....	16	69	225	118	428
White Shoal.....	44	0	10	52	106
Total oyster area.....	468	297	489	538	1,792
Total barren bottom.....					3,723
Total Baylor survey.....					5,515

As shown above, the barren bottom is equal to about 68 per cent of the area included within the Baylor lines, while the depleted area, which is almost uniformly worthless in its present condition, is equal to about 10 per cent. Assuming, as has been done in the discussion of the preceding grounds, that the bottom bearing a very scattering growth, of little or no present value so far as its market-oyster content is concerned, is capable of regeneration under the operation of natural agencies, the total present or prospective productive bottom constitutes about 22 per cent of the entire area. The following table shows the estimated present market-oyster content of the several rocks and their respective subdivisions according to density of growth:

CONTENT OF MARKET OYSTERS, PUBLIC GROUND No. 1 WARWICK COUNTY, BELOW DEEP CREEK.

Name of rock.	Dense.	Scatter- ing.	Very scatter- ing.	Depleted.	Total.
	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Browns Shoal.....	12,444	2,376	1,053	904	16,777
Gun.....	912	480	1,392
Kettle Hole.....	27,090	5,412	4,662	37,164
Thomas Point.....	8,745	8,378	5,100	2,667	24,890
Blunt Point.....	2,576	4,140	10,125	1,652	18,493
White Shoal.....	5,588	300	624	6,572
Total.....	57,355	20,306	21,780	5,847	105,288

The total content is nearly equal to that of the two grounds first described, but it will be observed that it is differently distributed, the dense areas bearing about twice the quantity of marketable oysters, the scattering about half as many, the very scattering about one-third more, and the depleted about three-fourths the quantity. With the exception of the depleted bottom, the average growth per acre is in each case somewhat greater than upon the grounds on the opposite side of the James and in the Nansemond River.

As will be understood from what has gone before, this distribution of the total content is to the distinct advantage of the oysterman, as a larger proportion of the oysters may be removed before work on the beds becomes unremunerative. The estimated available content of the beds embraced within this part of the public grounds—that is, the probable maximum yield during the present season—is shown in the following table:

AVAILABLE CONTENT OF MARKET OYSTERS, PUBLIC GROUND No. 1 WARWICK COUNTY, BELOW DEEP CREEK.

Name of rock.	Dense.	Scatter- ing.	Very scatter- ing.	Total.
	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Browns Shoal.....	9,800	1,200	250	11,250
Gun.....	600	600
Kettle Hole.....	18,900	2,700	800	22,400
Thomas Point.....	6,000	4,200	500	10,700
Blunt Point.....	2,000	2,000	2,000	6,000
White Shoal.....	4,000	4,000
Total.....	41,300	10,100	3,550	54,950

Practically four-fifths of the available oysters are found on the areas charted as bearing a dense growth, and about two-thirds of the remainder are on the areas of scattering growth. The bottoms covered by oysters in very scattering growths are slightly more productive than the average of the grounds previously described, but there are comparatively few spots on which a tonger could make a minimum livelihood. It is estimated that on the dense areas as a whole nearly three-fourths of the total content, on the scattering areas

about one-half, and on the very scattering areas not over one-sixth could be removed with profit. The depleted bottoms are worthless for their present product of market oysters.

The yield per acre of bottom included within the Baylor lines is considerably greater than on the grounds previously described, the average being almost 10 bushels, 2 bushels less than the average of the entire public area of Virginia in 1904, and considerably more than the average reported by the Census Bureau in 1908. The average available product of the oyster rocks, excluding all barren bottom but not that which is depleted, is about 30 bushels per acre. The average of the dense area is about 88 bushels, of the scattering area 34 bushels, and of the very scattering growth about 7 bushels per acre. The rocks in this ground are, on the whole, so far as present productiveness is concerned, in better condition than those across the river. The probable future productiveness of the beds, so far as the present existence of young oysters is concerned, is illustrated in the following table:

TOTAL CONTENT OF YOUNG OR SEED OYSTERS, PUBLIC GROUND NO. 1 WARWICK COUNTY, BELOW DEEP CREEK.

Name of rock.	Dense.	Scatter- ing.	Very scatter- ing.	Depleted.	Total.
	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Browns Shoal	8,568	6,248	2,376	1,130	18,322
Gun	1,188		992		2,180
Kettle Hole	81,786	13,662	19,980		115,428
Thomas Point	12,768	20,960	10,300	10,160	53,288
Blunt Point	2,736	13,317	23,625	4,956	44,634
White Shoal	14,728		1,080	2,756	18,564
Total	121,774	53,287	58,353	19,002	252,416

The exhibit here is much more favorable than on the rocks included in the grounds previously described, the average growth of young oysters on the dense and very scattering areas being over double that on the beds across the river, while that on the scattering area is about 35 per cent greater.

Practically everywhere on the areas of dense and scattered growth there is a prolific growth of young oysters and an abundance of clean shells, and there is no present prospect of the failure of these areas to continue to produce marketable oysters under ordinary conditions and with a reasonable enforcement of the laws. On the areas of very scattering growth the conditions are mixed, some places being well insured against the future and others being decidedly deficient in both young growth and clean shells. On Browns Shoal Rocks there is, with the exception of a few places, an abundance of shells; but there are only two or three patches where there is a supply of small oysters ample to replace the present market growth. On Kettle Hole Rock the conditions are good practically everywhere, but on Thomas Point and

Blunt Point Rocks there are but a few places where the young growth is prolific, and there are some in which neither young nor shells are found in even the minimum quantity requisite. On White Shoal Rock the condition on the areas of very scattering growth is in general satisfactory.

On the depleted bottoms as a whole the average growth of young oysters is about equal to that on similar bottom across the river, and there appears to be but little prospect of the future improvement of these areas, although there are a few spots on Thomas Point and White Shoal Rocks, in proximity to productive areas, where the growth of young is good.

MINOR PUBLIC GROUNDS.

In the Nansemond River and on the right side of James River there are several small public grounds, all of which are insignificant both in area and productiveness, and some of which were examined not at all or unsatisfactorily. They are as follows (somewhat more detailed data concerning some of them may be found in the descriptions of the individual rocks):

Nansemond County Ground No. 3. This was intended to include Holland Rock and at present contains in depleted bottom about 22 acres, on which there are a very few oysters and shells and about 33.9 acres of barren bottom.

Isle of Wight County Ground No. 2 contains about 9 acres of bottom of various degrees of productiveness, 24 acres of depleted and 16.8 acres of barren bottom. Its general condition is related in the description of Aaron Shoal Rock, its only natural bed. Isle of Wight County Ground No. 3 adjoins the preceding and has an area of $6\frac{1}{2}$ acres. It was not examined in the present survey. Isle of Wight County Ground No. 4 lies inshore of the preceding and covers about 3 acres of apparently depleted bottom.

Isle of Wight County Ground No. 5 embraces Creek Channel Shoal Rock, covering about 2 acres of depleted and 5.1 acres of barren bottom. Its present condition is described under the name of the rock.

PUBLIC GROUND NO. 1 WARWICK COUNTY, ABOVE DEEP CREEK.

The lower part of this ground, lying below Deep Creek, is within the area from which market oysters only can be taken and is therefore subject to the operations of the cull law. Its beds have been discussed in the foregoing. Above Deep Creek and Days Point, on both sides of the river, the cull law is suspended so far as young oysters are concerned, and, while shells must be returned to the beds, there is no limit on the minimum size of oysters which may be taken, the whole area being set apart for the production of seed for replanting.

It is necessary, therefore, in the discussion of the productiveness of this part of the James River oyster grounds, to adopt a different standard of productiveness. The whole oyster product of whatever size is involved in the question of the present value of the beds, whereas in the areas previously discussed the market oysters only could be considered, and the quantity of young was of interest merely as indicating the probability of the beds being maintained or repleted. In the discussion which follows here the maximum potential yield is considered as the production in excess of that which will give the tonger 4 bushels of oysters per day of tonging, not taking into consideration the time employed in culling out the shells and returning them to the beds.

This part of Ground No. 1 includes all oyster rocks on the left bank of the James River, from the mouth of Warwick River to the upper limit of oyster growth, near Deepwater Shoals Light-House. The following is a summary of the extent of the several rocks and the barren bottoms embraced within the Baylor lines:

AREAS OF OYSTER GROWTH, PUBLIC GROUND NO. 1 WARWICK COUNTY, ABOVE DEEP CREEK.

Name of rock.	Dense.	Scatter- ing.	Very scatter- ing.	Depleted.	Total.
	<i>Acres.</i>	<i>Acres.</i>	<i>Acres.</i>	<i>Acres.</i>	<i>Acres.</i>
Jail Island.....	227	198	14	508	947
Wreck Shoal.....	586	0	0	0	586
Dry Shoals.....	126	18	9	21	174
Point of Shoals.....	254	155	239	142	790
Swash.....	146	0	0	115	261
Mulberry Swash.....	422	34	20	29	505
Marshy Island.....	197	322	235	387	1,141
Long Shoal.....	331	10	84	79	504
V Rock.....	240	0	73	73	386
Moores.....	37	6	0	0	43
Horsehead.....	33	192	139	16	380
Deepwater Shoals.....	17	0	21	241	279
Total oyster area.....	2,616	935	834	1,611	5,996
Total barren area.....					6,896.8
Total Baylor survey.....					12,892.8

It will be noticed at once that the proportion of barren bottom to that actually included in the rocks as determined by the survey is somewhat smaller than in the grounds previously discussed, constituting about 53 per cent of the total. The depleted bottom, which, with practically no exceptions, is at present and potentially valueless, covers an additional 12 or 13 per cent, so that, assuming all the rest to be at present productive or capable of becoming so in the future, the oyster bottom covers about 35 per cent of the whole.

The following table exhibits the estimated total content of the several rocks and their subdivisions at the opening of the oyster season on September 15, 1909:

TOTAL CONTENT OF OYSTERS, PUBLIC GROUND NO. 1 WARWICK COUNTY, ABOVE DEEP CREEK.

Name of rock.	Dense.	Scatter- ing.	Very scat- tering.	Depleted.	Total.
	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Jail Island.....	32,461	21,582	392	4,064	58,499
Wreck Shoal.....	185,176	185,176
Dry Shoals.....	30,766	2,232	765	420	34,183
Point of Shoals.....	50,800	14,415	10,038	2,130	77,383
Swash.....	42,778	1,725	44,503
Mulberry Swash.....	127,444	3,604	2,600	1,247	134,895
Marshy Island.....	45,507	41,538	19,975	6,966	113,986
Long Shoal.....	79,771	640	5,040	1,264	86,715
V Rock.....	54,480	6,132	1,095	61,707
Moores.....	9,805	1,008	10,813
Horsehead.....	7,359	19,968	15,568	42,895
Deepwater Shoals.....	2,193	1,097	2,892	6,182
Total.....	668,540	104,987	61,007	21,803	856,937

It will be seen that the great preponderance of oyster production is on the dense areas, which exceed the bottoms of other character not only in their average productiveness but in their total area. The bottoms with a scattering growth, which in extent exceed the next lower grade by about 12 per cent, excel them in their total content by about 70 per cent, and are considerably more important in total production than the combined areas of very scattering oysters and depletion. Summarizing, the dense areas bear 78 per cent of the total content of the rocks, the scattering areas about 12 per cent, the very scattering about 7 per cent, and the depleted bottom about 3 per cent. Basing the computation on the basis previously defined and the data presented in the preceding two tables, we find the estimated maximum available product of the several rocks and their subdivisions to be as follows:

AVAILABLE CONTENT OF OYSTERS, PUBLIC GROUND NO. 1 WARWICK COUNTY, ABOVE DEEP CREEK.

Name of rock.	Dense.	Scatter- ing.	Very scat- tering.	Total.
	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Jail Island.....	25,000	12,000	100	37,100
Wreck Shoal.....	160,000	160,000
Dry Shoals.....	26,000	1,300	27,300
Point of Shoals.....	42,000	8,500	1,000	51,500
Swash.....	38,000	38,000
Mulberry Swash.....	116,000	1,800	1,000	118,800
Marshy Island.....	35,000	25,000	7,500	67,500
Long Shoal.....	68,000	300	1,500	69,800
V Rock.....	44,000	2,500	46,500
Moores.....	7,500	500	8,000
Horsehead.....	6,000	11,000	7,000	24,000
Deepwater Shoal.....	1,000	400	2,000
Total.....	569,100	59,900	21,500	650,500

The foregoing may be assumed to be the maximum quantity of seed oysters that can be profitably taken from the beds during the present season and the actual yield will probably fall considerably below the total exhibited in the table. Of the total, the areas of dense growth are capable of producing 88 per cent, of scattering growth 9 per cent, and of very scattering growth 3 per cent. The estimated yield per acre of bottom included within the boundary lines of this part of the bed is about 50 bushels. This low average of production is of course induced by the large area of barren and depleted or practically barren bottom included in the Baylor lines. If we compare the average of the whole area with that of the best bottom in the natural rocks under discussion the paucity of the former is equally impressive, the dense areas of the region under discussion having an average total content of about 256 bushels per acre and a promised yield during the present season of 213 bushels, over four times the average of the beds as a whole. The average available product of the areas of scattering growth is about 64 bushels per acre, and of very scattering growth about 13 bushels, both yields being far below what they should produce under proper conditions.

Upon the dense areas as a whole the present production and the promise for the future are both good, and on the area of scattering growth, while the present production is fair, the quantity of shells is such as to promise a better yield in the future, should there come a season of heavy and general strike.

On the bottoms rated as bearing a very scattering growth the conditions as a whole are not such as to yield much profit to the tonger, though in some places he could make a living wage for a short period. In most places on bottom of this character the quantity of clean shells is such as to give indifferent prospect of the future regeneration of the beds.

The depleted bottom, excepting in a few places near Deepwater Shoals Light-House, bears shells in such small quantities as to make exceedingly remote the probability of any material improvement under natural conditions.

PUBLIC GROUND NO. 1 ISLE OF WIGHT COUNTY.

This ground extends as a narrow strip along the right bank of James River from close to the shore line out to the main channel, between Rock Wharf and Days Point Shoal. It lies wholly within the area set apart for seed production, and the statements in regard to the methods employed in computing the productiveness of the several parts of the preceding ground are applicable to this as well.

Compared with the extensive areas occupied by the rocks across the river in Warwick County, the beds included in this ground are

insignificant. For the purposes of this report it is considered to include four natural rocks, although the largest of these, for which no name was obtained from the oystermen, may be locally recognized by names for its constituents severally. The general condition and extent of the bed are shown in the following table:

AREAS OF OYSTER GROWTH, PUBLIC GROUND No. 1 ISLE OF WIGHT COUNTY.

Name of oyster rock.	Dense.	Scatter- ing.	Very scat- tering.	Depleted.	Total.
	<i>Acres.</i>	<i>Acres.</i>	<i>Acres.</i>	<i>Acres.</i>	<i>Acres.</i>
Rock Wharf Shoals.....	18	0	0	8	26
Between Rock Wharf and Spindle Rock.....	45	0	5	37	87
Spindle.....	14	0	3	2	19
Days Point Shoal.....	4	0	0	0	4
Total oyster area.....	81	0	8	47	136
Total barren area.....					589
Total Baylor survey.....					725

The area of barren bottom as compared with the extent of the ground is relatively large, constituting about 81 per cent, and the depleted bottom, which is at present worthless and holds forth no promise of improvement, adds an additional 7 per cent to the wholly unproductive bottom. The area of dense growth, which is undoubtedly productive, covers about 11 per cent of the whole, while the bottom bearing very scattered oysters, which is at present practically incapable of yielding a living wage to the tonger, covers about 1 per cent.

The following table shows the estimated total content of oysters on the rocks at the end of August, 1909:

TOTAL CONTENT OF OYSTERS, PUBLIC GROUND No. 1 ISLE OF WIGHT COUNTY.

Name of rock.	Dense.	Scatter- ing.	Very scat- tering.	Depleted.	Total.
	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Rock Wharf Shoals.....	2,520			88	2,608
Between Rock Wharf Shoals and Spindle Rocks.....	6,300		230	666	7,196
Spindle.....	1,960		81	24	2,065
Days Point Shoal.....	664				664
Total.....	11,444		311	778	12,533

The total content of the ground as a whole averages about 17 bushels per acre. Practically all of this is borne by the small fraction of the bottom classed as dense, on which the average production is at the rate of about 141 bushels per acre, considerably less than on the areas of dense growth on the great beds across the channel.

The estimated available content on these beds—that is, the quantity which may be removed before tonging will cease to pay even a very small assumed minimum livelihood—is as follows:

AVAILABLE CONTENT OF OYSTERS. PUBLIC GROUND NO. 1 ISLE OF WIGHT COUNTY.

Name of rock.	Dense.	Scatter- ing.	Very scatter- ing.	Total.
	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Rock Wharf Shoals.....	2,000			2,000
Between Rock Wharf Shoals and Spindle Rock.....	5,000		50	5,050
Spindle.....	1,500		20	1,520
Days Point.....	500			500
Total.....	9,000		70	9,070

Practically all of the available supply of oysters on this ground is therefore on the bottom classed as dense and the area of very scattering growth is negligible. The available product is of the average density of 111 bushels per acre. On the dense area the shells are sufficient, on the very scattering area they are in fair quantity, while on the depleted ground they are deficient.

SUMMARY.

The public grounds in the region covered by the survey and of which a detailed discussion is found in the preceding pages cover an area of 26,408.4 acres as computed in the report of the Baylor survey. Of this acreage, 12,790.6 acres lie below the line drawn between Deep Creek and Days Point and 13,617.8 acres lie above that line. The beds of the former region are available for the production of marketable oysters only, the law requiring that all oysters under 3 inches long be returned to the beds, while the latter region is set apart for the production of seed oysters, and the cull law is not applicable except in so far as it forbids the removal of shells.

Of the entire area the recent survey shows that 3,227 acres may be classed as bearing a dense growth, 2,078 as scattering, 1,848 as very scattering, 3,884 as depleted, and 15,371.4 as barren. The barren and depleted bottoms together comprise 19,255.4 acres, or about 73 per cent of the total, and all bottom which can be construed as productive aggregates 7,153 acres, or 27 per cent of the entire bottom included within the Baylor lines. Owing to the difference in the provisions of the law applicable to the two regions and the resultant difference in the character of their product, it is necessary to present separate summaries of their present condition.

MARKET OYSTER AREA.

The beds of this region are shown on chart 1 accompanying this report, to which, and to the preceding pages, readers are referred

for detailed data. The following table summarizes the extent and character of the bottom included within the Baylor lines:

SUMMARIZED STATEMENT OF MARKET OYSTER AREAS ON PUBLIC GROUNDS.

Name of ground.	Dense.	Scatter- ing.	Very scat- tering.	Depleted	Barren.	Total.
	<i>Acres.</i>	<i>Acres.</i>	<i>Acres.</i>	<i>Acres.</i>	<i>Acres.</i>	<i>Acres.</i>
Nansemond No. 2.....	} 256	646	514	1,640	4,106.8	7,162.8
Isle of Wight No. 6.....		0	0	22	33.9	55.9
Nansemond No. 3.....		2(?)	4	3	24	16.8
Isle of Wight No. 2.....						
Isle of Wight No. 3.....						
Isle of Wight No. 4.....						
Isle of Wight No. 5.....	0	0	0	2	5.1	7.1
Warwick No. 1 and No. 2 (below Deep Creek).....	468	297	489	538	3,723.0	5,515.0
Total.....	726	947	1,006	2,226	7,885.6	12,790.6
Per cent.....	5.7	7.4	7.9	17.4	61.6	100.0

It is estimated that the bottoms embraced by the several grounds, classified in accordance with their relative productiveness, have a total content of market oysters as follows:

SUMMARIZED CONTENT OF MARKET OYSTERS ON PUBLIC GROUNDS.

Name of ground.	Dense.	Scatter- ing.	Very scat- tering.	Depleted	Total.
	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Nansemond No. 2.....	} 28,468	39,087	17,855	21,052	106,462
Isle of Wight No. 6.....					
Nansemond No. 3.....		200	132	69	330
Isle of Wight No. 2.....				132	533
Isle of Wight No. 3.....					
Isle of Wight No. 4.....					
Isle of Wight No. 5.....				22	22
Warwick No. 1 and No. 2 (below Deep Creek).....	57,355	20,306	21,780	5,847	105,288
Total market oysters.....	86,023	59,525	39,704	27,383	212,635
Average per acre.....	118	63	39	12	
Per cent of total.....	40.3	28.0	18.7	13.0	100

This table is more or less misleading, as the real factor involved is the quantity of oysters which can be profitably removed from the beds. It must be obvious that the total quantity lying on the bottom can not be regarded as commercially available, for when the density of growth is reduced below a more or less definite minimum the value of the average catch will fall below a minimum living wage and work will cease. The minimum average quantity per unit of bottom which will suffice to support commercial operations will depend upon the price of oysters and the depth of water. In this report the price is placed at 45 cents per bushel, and although it will vary somewhat on the different beds and at different times, it is not practicable to make distinctions. The price adopted is based on the testimony of a number of oystermen as to their returns in recent years. The depth of water is a highly variable factor, and as it is of prime importance in

computing the availability of the oysters lying on the bottom, its variations have been given the fullest possible consideration. For a discussion of the general principles on which the quantity of oysters available with profit have been determined, the reader is referred to preceding pages. For the market oyster beds as a whole the following table gives a summary:

SUMMARY OF AVAILABLE CONTENT OF MARKET OYSTERS ON PUBLIC GROUNDS.

Name of ground.	Dense.	Scatter- ing.	Very scat- tering.	Total.
	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Nansemond No. 2.....	21,150	20,325	1,600	43,075
Isle of Wight No. 6.....				
Nansemond No. 3.....	150	50		200
Isle of Wight No. 2.....				
Isle of Wight No. 3.....				
Isle of Wight No. 4.....				
Isle of Wight No. 5.....				
Warwick No. 1 and No. 2 (below Deep Creek).....	41,300	10,100	3,550	54,950
Total available market oysters.....	62,600	30,475	5,150	98,225
Average per acre.....	86	32	5	
Per cent.....	63.7	31.0	5.3	100.0

This may be regarded as a maximum estimate of the probable yield of the beds during the season of 1909-10. Owing to the low basis adopted as a minimum wage the yield may not reach the quantity indicated, as it is doubtful whether the beds can be profitably fished as closely as has been assumed. A yield of \$1.35 per full day of tonging will leave a very small balance after culling and other expenses are deducted, and the beds undoubtedly will be abandoned for the season before this degree of depletion has been reached. For this reason the only parts of the natural rock which can be classed as really productive are those designated as dense and scattering, which furnish, according to the foregoing estimates, about 95 per cent of the available product while constituting only about 13 per cent of the total area of the public grounds under consideration.

Taken as a whole, though there are exceptions noted in the preceding accounts of the individual rocks, the areas covered with very scattering growths are of but little present value, their total estimated available product during the present season being valued at less than \$2,500, or about \$2.50 per acre. There is, however, another phase to the question which has been touched on in the more detailed accounts of the individual rocks. This is the possibility of future improvement, and is dependent upon the existing quantity of young oysters and the presence of an ample supply of clean shells to serve as places of attachment for future generations of young. The quantity of young oysters less than 3 inches long on the public grounds under discussion at the opening of the present oyster season was as follows:

SUMMARIZED CONTENT OF YOUNG OYSTERS ON PUBLIC GROUNDS.

Name of ground.	Dense.	Scatter- ing.	Very scat- tering.	Depleted.	Total.
	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>	<i>Bushels.</i>
Nansemond No. 2.....	31, 818	86, 050	26, 854	57, 038	201, 760
Isle of Wight No. 6.....					
Nansemond No. 3.....				50	50
Isle of Wight No. 2.....	250	500		2, 500	3, 250
Isle of Wight No. 3.....					
Isle of Wight No. 4.....					
Isle of Wight No. 5.....					
Isle of Wight No. 6.....					
Warwick No. 1 and No. 2 (below Deep Creek).....	121, 774	53, 287	58, 353	19, 002	252, 416
Total.....	153, 842	139, 837	85, 207	78, 590	457, 476
Average per acre.....	212	148	84	35	
Per cent.....	33. 7	30. 6	18. 6	17. 1	100. 0

Here again is evidence that the areas of dense and scattering growth should be regarded as not only at present but prospectively productive beds. In quantity, and to a greater extent numerically, the young are considerably in excess of the market oysters, and, as in both classes of bottom under consideration the latter are sufficient to render the bottom undoubtedly at present productive, the abundance of young is sufficient to continue productiveness, under proper regulations as to culling, for at least two years. After the lapse of that period the condition will depend upon the extent of the strike, and other factors concerning which nothing can be predicted.

So far as the areas covered with a very scattering growth are concerned closer scrutiny is required. As these bottoms are on the verge of depletion in respect to market oysters, the proportion of young to large oysters should be greater to insure that the conditions will improve in the future.

The writer is not in possession of definite experimental data applicable specifically to the James River, but from a knowledge of conditions in other parts of the Chesapeake region he feels justified in assuming that oysters as an average will become fit for market in from two to three years from the time of fixation or setting. If experience elsewhere be a guide, some oysters will grow more rapidly and some less rapidly; but two years may be adopted, with very little question, as an irreducible minimum for the average age at which they can be advantageously put on the market. On this assumption and neglecting, for the time being, the question of mortality, it is at once apparent that to maintain the present status there must be two small oysters for each market oyster killed or caught.

There is no way to determine, without long and painstaking observations, the actual average mortality at various ages on the natural rocks of James River. The experience of planters of seed oysters is valueless in this connection, being based on oysters handled and

otherwise subjected to abnormal conditions. The various locations of the natural-bed oysters and the consequently variable conditions to which they are exposed introduce a factor for which it is difficult to make allowances, and it seems impossible to do more than hazard a guess as to the proportion of young oysters now on the beds which will die before becoming marketable. It is probable that it will be somewhere between 25 and 50 per cent. Considering the size of young oysters found by the survey, the mortality may be less than the former and excepting under unfavorable conditions can hardly be greater than the latter. Assuming that 25 per cent of the young now on the beds will die before reaching a marketable size, there should be on the beds, in order to maintain their present condition, 2.66 young for each marketable oyster removed. If the loss be assumed at 50 per cent there should be 4 young per market oyster.

The following table exhibits the actual average numerical proportion of young oysters to marketable found on the several beds:

NUMERICAL PROPORTION OF YOUNG GROWTH AND MARKET OYSTERS IN THE MARKET-OYSTER AREA ON BOTTOMS BEARING VERY SCATTERING GROWTH.

Name of rock.	Oysters less than 1 inch long.	Oysters between 1 and 3 inches.	Total.
Nansemond Ridge.....	0.42	1.82	2.24
Drum Shoal.....	.62	3.29	3.91
Newport News.....	.50	5.27	5.77
Cruiser Shoal.....	.55	3.83	4.38
Flat Rock, etc.....	.69	2.58	3.27
High Shoal.....	1.81	2.47	4.28
Trout Shoal.....	.55	6.81	7.36
Dog Shoal.....	.27	1.13	1.40
Fishing Point.....	1.11	1.94	3.05
Between Fishing Point and Ballards Marsh.....	.06	.00	.06
Ballards Marsh.....	3.50	11.40	14.90
Aaron Shoal.....	.66	8.00	8.66
Browns Shoal.....	.80	4.00	4.80
Gun.....	.19	4.28	4.47
Kettle Bottom.....	.63	8.27	8.90
Thomas Point.....	.21	4.06	4.27
Blunt Point.....	.35	4.56	4.91
White Shoal.....	.87	5.36	6.23
Average.....	.76	4.39	5.15

It will be observed that, on the assumption of the smaller death rate, Nansemond Ridge, Dog Shoal, and the small beds near Ballards Marsh are the only rocks which appear to lack sufficient young growth on the very scattering areas to maintain them in their present condition. Assuming the higher rate of mortality, Drum Shoal, the small bed near Flat Rock, and Fishing Point Rocks must be added to the list, though when we consider that many of the market oysters now on the bottom can not be taken with profit, it would appear that even these rocks are capable of improving under a rigid observance of the cull law. The other rocks, under either assumption as to mortality, probably bear a sufficient number of young to

maintain their present condition or to cause slight improvement in their productiveness, and the real question at issue largely resolves itself into a matter of their present productiveness, which has been already discussed. Most of these bottoms are sufficiently clothed with shells to insure their share of a good strike. The depleted areas, excepting in the places specifically mentioned in the descriptions of the individual rocks, may be regarded as hopeless of recuperation under natural conditions.

The barren bottoms, which preceding tables show to constitute a very large proportion of the areas of the public beds, are in many cases so situated as to be of necessity and for practical considerations impossible of separation from the natural beds without injury to the future of the latter or without due regard to the question of policing and administration. There are, however, certain large areas readily separable from the public grounds, and the latter would suffer practically no diminution in really productive bottom as a result of the severance.

SEED-OYSTER AREA.

This region lies above the line drawn between Deep Creek and Days Point, and is shown on chart 2 accompanying this report. The following table summarizes the extent and condition of the bottoms of different degrees of productiveness included within this part of the Baylor survey:

SUMMARIZED STATEMENT OF OYSTER GROWTH ON SEED AREAS.

Name of ground.	Dense.	Scatter- ing.	Very scat- tering.	Depleted.	Barren.	Total.
Warwick No. 1 (above Deep Creek).....	<i>Acres.</i> 2,420	<i>Acres.</i> 1,131	<i>Acres.</i> 834	<i>Acres.</i> 1,611	<i>Acres.</i> 6,896.8	<i>Acres.</i> 12,892.8
Isle of Wight No. 1.....	81	0	8	47	589.0	725.0
Total.....	2,501	1,131	842	1,658	7,485.8	13,617.8
Per cent.....	18.4	8.3	6.2	12.2	54.9	100.0

The following table furnishes an estimate of the total content of seed oysters present on the bottoms of varying productiveness at the beginning of the oyster season on September 15, 1909:

SUMMARIZED CONTENT OF OYSTERS ON SEED AREAS.

Name of ground.	Dense.	Scatter- ing.	Very scat- tering.	Depleted.	Total.
Warwick No. 1 (above Deep Creek).....	<i>Bushels.</i> 668,540	<i>Bushels.</i> 104,987	<i>Bushels.</i> 61,607	<i>Bushels.</i> 21,803	<i>Bushels.</i> 856,937
Isle of Wight No. 1.....	11,444	311	778	12,533
Total.....	679,984	104,987	61,918	22,581	869,470
Average per acre.....	272	93	73	13
Per cent.....	78.2	12.1	7.1	2.6	100.0

A considerable proportion of these oysters could not be profitably removed from the beds, being either too sparsely distributed in the first place or constituting a necessary remnant which would become too scattered after tonging had been carried on for a period on bottoms of greater initial productiveness. To show the estimated maximum possible yield of the beds during the present season the following table has been prepared, covering the entire area of seed beds in the James River:

SUMMARY OF AVAILABLE CONTENT OF OYSTERS ON SEED AREAS.

Name of ground.	Dense.	Scatter- ing.	Very scatter- ing.	Total.
Warwick No. 1 (above Deep Creek).....	<i>Bushels.</i> 569,100	<i>Bushels.</i> 59,900	<i>Bushels.</i> 21,500	<i>Bushels.</i> 650,500
Isle of Wight No. 1.....	9,000	70	9,070
Total.....	578,100	59,900	21,570	659,570
Average per acre.....	232	53	25
Per cent.....	87.7	9.1	3.2	100.0

In preparing the data on which this table is based it has been assumed that the seed will bring 30 cents per bushel and that no bottom can be considered productive when its yield is reduced below 4 bushels per day of actual tonging, excluding the time occupied in culling. As in the preceding pages of this report, the probable yield is based on the density of the oyster growth and the depth of water on the several parts of each bed.

Although the data employed differs somewhat from that used in the discussion of the bottoms below Deep Creek, owing to the lower price brought by seed as compared with market oysters, the financial return to the tonger from the bottoms designated as respectively dense, scattering, and very scattering is essentially the same. The minimum yield assumed to place a given area above the grade of depleted bottom is valued at \$1.20 per day at the prices recently prevailing, and this can not be regarded as other than an extreme minimum, because, when the number of idle days is taken into consideration, a tonger could not afford to work for such low wages. The limit is justifiable only in consideration of the fact that before the dense and scattering areas are reduced to a level so low they will have yielded to the tonger an average daily wage much in excess of this.

If the price of oysters falls below 30 cents per bushel, it will not be profitable to work the beds so closely as was contemplated in the preparation of the above table. At the prices reported as current on the James River in November, 1909, namely, 20 cents per bushel for seed, the estimated catch on the area of very scattering growth may be eliminated, that on the scattering bottom reduced by at least 50 per cent and on the dense areas by about 15 per cent, lowering the

total estimated catch to about 500,000 bushels as a maximum which could be taken with profit.

As was the case with the market-oyster beds, the areas described as bearing dense and scattering growths may be dismissed from further discussion as being at present productive. The areas bearing a very scattering growth are debatable, with seed oysters selling at 30 cents per bushel, but would be undoubtedly entirely neglected by the tongers were the price to fall to 20 cents. Their estimated yield at the former price is about \$7.50 per acre, and from the entire area of 842 acres the total product during the present season would not exceed in value \$6,500, even if the tongers were willing to work for an average of about \$1 per day, exclusive of the time lost through bad weather.

The future of these areas of very scattering growth is difficult to forecast. When, as in the area under discussion, there is not and from the nature of the case should not be any application of the cull law, there is no young growth which can be pointed to as coming forward to replace the larger oysters removed. Young and old alike are taken and the only oysters left are the residuum which it is unprofitable to take. In other words, the annual increment is taken or may be taken in the months immediately following its deposit. The health and perpetuity of the beds depend upon the quantity of clean shells exposed on the bottom ready for the strike which each season may produce. Over the very scattering bottoms of this part of James River there is a fair quantity of shells and under the proper conditions these areas may become more productive.

The depleted bottoms, as a whole, have neither present nor prospective value under natural conditions, though the bottom is generally of such character that if it were feasible to rent it for purposes of oyster culture it could be made highly productive. Much of it is so situated, however, that it is debatable whether, for reasons of administration, it would be advisable to alienate it from the public grounds. An inspection of the charts will show that, excepting along the shores, these bottoms are generally in the midst of productive areas. Concerning the great area of barren bottoms the same statement holds true in part, a considerable proportion of it lying in the channels and deeper holes between the beds or in other situations which would make it difficult to delimit it from the public grounds in a manner to facilitate the policing of the public rocks and prevent abuses which experience shows would undoubtedly be attempted.

There are, however, certain areas in considerable blocks which could be set apart for purposes of oyster culture without materially reducing the area of the natural rocks included in the public grounds. These places can be determined by an inspection of the chart.

CONCLUSION.

The foregoing gives, in detail and summary, the facts as to the condition of the oyster beds of James and Nansemond rivers immediately prior to the opening of the present oyster season, the period at which the beds are at their maximum apparent productiveness. Within a few weeks, under the intensive fishery which they sustain, the quantity of oysters on the beds will be vastly reduced and long before the close of the season they will become so impaired that work on them will be practically abandoned for the time being. In other words, it is for a part of the season only that these beds will offer a livelihood to the tongers, who for the rest of the year must seek a living either in the employ of the oyster grower or in some other occupation not connected with oyster fishing.

In the determination of the nature of tidal bottoms, with respect to their being regarded as oyster rock or barren bottom, the prime consideration is whether they will afford, either at present or prospectively, a sufficient quantity of oysters to provide a livelihood to those who work on them. It is manifest that a few oysters which could never be taken with profit should not entitle the bottom on which they lie to be regarded as an oyster bed within the meaning of the laws. To so regard them would be contrary to common sense, economic principles, and judicial decisions.

The author has avoided a definition of what constitutes a livelihood, believing that to be a matter which is more properly for determination by the state authorities should its definition become necessary for purposes of legislation or administration. In the preparation of the foregoing report, however, it has been necessary to adopt some standard for the classification of the various densities of oyster growth in the several beds, and for purposes of convenience the limit between the bottoms regarded as depleted and those of the lowest class of productiveness has been placed at a minimum believed to be reasonably irreducible. The subdivisions of productiveness differ by such small quantities that should it appear that the lowest is too low the next higher can be regarded as the minimum without impairing the value of the data adduced in the report, though, as is elsewhere indicated, this would dictate a reduction in the estimated total available product of oysters for the season.

Under the terms of the resolution of the State Board of Fisheries, which was made the basis of the request for the survey preferred to the Bureau of Fisheries by the Governor of Virginia, the author is not warranted in offering recommendations as to the use which might be made of the facts developed in the preceding pages. It may not be inappropriate, however, to point out the several avenues of procedure which it is possible to follow in respect to the oyster

bottoms of the region discussed. These resolve themselves into three: (1) The maintenance of the integrity of the public grounds as now constituted; (2) their abolition in toto; and (3) a middle course which will preserve to the public the productive bottoms practically in their entirety while throwing open to oyster planting a large part of the barren and unproductive bottom now included within the public grounds. The principal arguments for and against these propositions may be epitomized as follows:

1. The first course—that the beds be retained in the present status—hardly needs discussion. It has been tried and its results are known, largely as the effect of the acrimonious disputes to which it has given rise. The matters of fact which have been at issue in these interminable discussions, as to whether or not the public grounds embraced any considerable area of barren bottom, have been dealt with in the preceding pages and speak for themselves. It should be pointed out, however, that while the barren area is shown to constitute a considerable proportion of the whole bottom, much of it is so related to the productive bottom that it could not be eliminated under any scheme permitting of practical administration.

2. The second alternative—the total abolition of the public grounds and its corollary, the opening of the whole area for leasing—is drastic. On broad economic grounds the proposition is as logical and legitimate as the sale of public timber land or the breaking up of the great public ranges of the West into holdings in severalty, and, as the oyster is sessile, it has nothing in common with an alienation of the common fishery for nomadic species. The law has already recognized that under conditions an oyster in situ may be property, while a wandering fish can not become such until caught. The breaking up of the public grounds into leaseholds under private control would increase their productiveness precisely as the breaking up of the common ranges of the West has resulted in economic efficiency and greater productiveness. This course would, furthermore, yield a return to the State, where there is now a net outlay for policing the public grounds, though this aspect of the matter is one which should always be held subservient to the major consideration—the welfare and prosperity of the citizen.

On the other side of the question it is necessary to consider the effect of so drastic an innovation upon the welfare of a large body of persons whose livelihood in part is at present dependent upon the situation to which the policy of the State has given the aspect of presumed permanency. Immediately upon the alienation of the public beds the men engaged on them for part of the season are, for the time being, forced from the category of independent workers into that of employees, unless they themselves elect to take up

bottoms in severalty. The value of this objection is mainly sentimental, but is not less real on that account.

3. The third course mentioned, the retention of the actually productive bottom for the use of the public and the opening of all barren bottom practicable for leasehold from the State, is essentially a compromise between the other two and presents fewer difficulties than either. The valid objections to it are mainly concerned with administration. By retaining the present natural beds intact the tongers would be left in possession of everything of value to which they now have access, while the opening of the barren bottoms for lease would make productive considerable acres now valueless to all. The tongers would still have the option of independent work on the natural rocks; they would have increased opportunities of employment by the planters; and some of them could themselves lease bottoms for their own use. In every way it would appear to be economically advantageous to the industry and the State.

In considering the subject, however, it should be borne in mind that, while this report shows a preponderance of barren bottom within the public grounds, much of it, owing to its location, is practically incapable of separation from the natural rocks. An inspection of the chart will show that many of the barren bottoms are between or in the midst of naturally productive bottoms. To exclude them would make necessary an undue multiplication of the public grounds, with an attendant difficulty in policing.

Effectually to prevent depredations on the natural rocks under the guise of work on adjoining planted grounds, which is a difficulty with which the oyster police will have to contend, the public areas should be as few and as compact as possible, and the boundary lines should be straight and easily defined. For this reason the public grounds to be established must, for very practical considerations, necessarily include a considerable proportion of barren bottom. Any readjustment of the lines of the Baylor survey should be based on reasonable compromise and adopted only after careful consideration by the State. It is believed that the foregoing descriptions and the accompanying charts will furnish a reliable basis for a revision, should the State deem it wise to undertake it.

DESCRIPTION OF CHARTS.

The public grounds are platted from the published sheets of the Baylor survey, and their boundaries are shown in broken black lines.

The depths, which are expressed in feet as referred to mean low water and the symbols designating the consistency of the bottom, are the characteristics selected from a large number of observations. The oyster beds are included within solid red lines, and the density of the oyster growth is indicated by the relative intensity of the shading, and is based on the quantity of culled oysters which can be taken by a tonger working nine hours per day, not including the time occupied in culling.

Chart 1 covers the area from which market oysters only may be removed, and the bases of the classifications of oyster growth are as follows: Dense, yielding over 8 bushels of market oysters per day's tonging; scattering, between 5 and 8 bushels; very scattering, between 3 and 5 bushels; depleted, under 3 bushels.

Chart 2 embraces the beds of the upper part of James River, on which the cull law is inoperative so far as it pertains to the size of the oysters, and which are therefore devoted to the production of seed oysters for replanting. The classification is as follows: Dense, yielding over 12 bushels of seed oysters per day's tonging; scattering, between 8 and 12 bushels; very scattering, between 4 and 8 bushels; depleted, under 4 bushels.

The unshaded areas within the boundaries of the public beds as charted represent barren bottom.

O







Boundaries of Public Beds shown in broken black lines.
 Natural Oyster Beds, shown in red:
 Area of dense growth of market oysters [diagonal lines] [diagonal lines]
 " " scattered " " " " [diagonal lines] [diagonal lines]
 " " very scattered " " " " [diagonal lines] [diagonal lines]
 " " depleted " " " " [diagonal lines] [diagonal lines]
 Stations at which dense growth of young occur [circle]

Soundings:
 Depth at mean low water expressed in feet.
 Bottom symbols: h, for hard; s, for soft.

Note: For more detailed explanation see text.




Department of Commerce and Labor
 BUREAU OF FISHERIES


Survey of
 THE PUBLIC OYSTER BEDS
 of
 JAMES RIVER VIRGINIA

By Fisheries Str. FISH HAWK, Chf. Debn W.Martin, U.S.N., Comdg
 Under the direction of
 H.F.Moore, Assistant, Bureau of Fisheries


JULY, AUGUST, SEPTEMBER
 1909



E. Veitch, Draftsman











Boundaries of Public Beds shown in broken black lines.
 Natural Oyster Beds, shown in red:
 Area of dense growth of seed oysters
 " " scattered " " "
 " " very scattered " " "
 " " depleted " " "
 Soundings:
 Depth at mean low water expressed in feet.
 Bottom symbols: h, for hard; s, for soft.
 Note: For more detailed explanation see text.


Department of Commerce and Labor
 BUREAU OF FISHERIES
 Survey of
THE PUBLIC OYSTER BEDS
 of
JAMES RIVER VIRGINIA

By Fisheries Str. FISU HAWK. Chf. Bos'n W. Martin, U.S.N., Comdg
 Under the direction of
 H.P. Moore, Assistant, Bureau of Fisheries
 JULY, AUGUST, SEPTEMBER
 1909













LIBRARY OF CONGRESS



0 002 896 853 A